Mon. Not. R. Astron. Soc. **000**, 1–28 (2012) Printed 11 January 2013 (MN LaTeX style file v2.2)

arXiv:1301.1975v1 [astro-ph.SR] 9 Jan 2013

The death of massive stars - II. Observational constraints on the progenitors of type Ibc supernovae

John J. Eldridge[1] *, Morgan Fraser[2], Stephen J. Smartt[2],
Justyn R. Maund[2], R. Mark Crockett[2]

[1] *The Department of Physics, The University of Auckland, Private Bag 92019, Auckland, New Zealand*
[2] *Astrophysics Research Center, School of Mathematics and Physics, Queen's University Belfast, Belfast BT7 1NN, UK*

11 January 2013

ABSTRACT

The progenitors of many core-collapse supernovae have now been identified directly on pre-discovery imaging of their host galaxies. These are all of type II and all of the most common sub-types have at least one detection (II-P, II-L, IIb and IIn). Here we present an extensive search for the progenitors of type Ibc supernovae in all available pre-discovery imaging since 1998. To date there have been 12 type Ibc supernovae with either deep ground-based imaging or Hubble Space Telescope archival imaging of their explosion sites. Our analysis of this data set shows there is no detection of any progenitor in these images, with the deepest limits probing absolute magnitudes of between -4^m and -5^m in broad-band B, V and R. We compare the magnitude limits with the observed Wolf-Rayet population in the Large Magellanic Cloud, which show a range in their broad-band absolute magnitudes between -2^m and -8^m. There is only a 15 per cent probability that this is actually the progenitor population and that we have failed to detect a Wolf-Rayet progenitor by chance selection. Comparing to the observed Wolf-Rayet populations in NGC5068 and NGC7793, we find an essentially zero probability that this population could give rise to the observed Ibc population. If SN Ibc progenitors are drawn from such populations then either Wolf-Rayet stars evolve significantly in the last stages of their lives, or we have underestimated circumstellar dust and extinction towards the progenitors. We review the relative rates of Ibc SNe, finding agreement with previous work that they occur too frequently to originate only from progenitors with initial masses $> 25\mathrm{M}_{\odot}$. The ejecta mass estimates from lightcurve modelling of Ibc SNe indicate typical ejecta masses between 1-3$\mathrm{M}_{\odot}$, which is somewhat low to be compatible with massive Wolf-Rayet progenitors. Hence we propose that a Wolf-Rayet origin for the bulk of the normal Ibc population is unlikely. We present results from binary stellar evolution calculations and population models that show that stars less than approximately 20$\mathrm{M}_{\odot}$, in interacting binaries can fit all of these observational constraints. These stars lose their hydrogen envelopes through either Roche-lobe overflow or common envelope evolution and end their lives as low mass helium stars. We show that such helium stars in model binaries retain a low mass hydrogen envelope until after the end of core helium burning and lose the last of the hydrogen approximately 10^4 years before core-collapse. Hence it is not surprising that they have been difficult to identify, but some viable Ibc progenitor systems exist.

Key words: stars: evolution – binaries: general – supernovae: general – stars: Wolf-Rayet – stars: supergiants

1 INTRODUCTION

Massive stars live fast and die young; core-collapse supernovae (SNe) are the dramatic explosions which mark their deaths. During these events a large amount of energy is de-

---

* E-mail: j.eldridge@auckland.ac.nz

© 2012 RAS

posited into the surrounding interstellar medium, while elements heavier than boron are created through explosive nucleosynthesis. The physical mechanism of core-collapse in a star with an iron core is well established as the main driving source behind the vast majority of core-collapse explosions (Janka 2012; Burrows 2012). However the structure of the star before explosion and the variety in explosions (kinetic energies, luminosities and variation in the production of radioactive ^{56}Ni) are still not well understood. The observational characteristics are critically affected by the evolutionary history of its progenitor star and uncertainties remain (most notably in convection, mass-loss and rotation) that limit our quantitative understanding (see Langer 2012, for a discussion of current limitations).

Observed SNe are first divided into Types I and II, based respectively on the absence or presence of hydrogen in the early-time optical spectrum (Filippenko 1997; Turatto, Beneti & Pastorello 2007). These two broad types are further divided into several subtypes. The most common Type II SNe are Type IIP, characterised by a long plateau phase in their lightcurve of constant luminosity, which lasts for several months. Type IIL SNe have a linear decay to their light curve, while Type IIb SNe only show hydrogen in their spectra for the first few weeks before it disappears. Type IIn SNe are slightly different, the hydrogen is seen as narrow (< 100 km s^{-1}) emission lines, which arise from interaction with dense, optically thick circumstellar material. Type Ia SNe are hydrogen and helium poor SNe that are believed to be thermonuclear explosions arising as a result of accretion onto a carbon-oxygen white dwarf and are not considered further here (see Hillebrandt & Niemeyer (2000) for a review). The other hydrogen-poor Type I SNe are subdivided into Types Ib and Ic, which either do or do not show helium in their spectra. The type Ibc SNe[1] have long been associated with Wolf-Rayet progenitors (e.g. Begelman & Sarazin 1986; Gaskell et al. 1986).

In Paper I of this series, Smartt et al. (2009), the progenitors of Type IIP SNe were identified as red supergiants (RSGs) with massive and extended hydrogen envelopes. This conclusion was based on an analysis of pre-explosion images (mostly from the *Hubble Space Telescope*) for twenty nearby SNe. These images were used to detect or place limits on the luminosity and thus mass of the progenitors. For the other subclasses of Type II supernovae there are some detections, but insufficient numbers to consider in detail as an ensemble. There have been three detected Type IIb progenitors for SNe 1993J, 2008ax and 2011dh, one transitional Type IIP/L progenitor for SN 2009kr and a progenitor detected for the peculiar Type II SN 1987A (Walborn et al. 1987; Aldering, Humphreys & Richmond 1994; Maund et al. 2004; Crockett et al. 2007; Elias-Rosa et al. 2010; Fraser et al. 2010b; Maund et al. 2011; Arcavi et al. 2011). Together these observations suggest that if the progenitors of Type II SNe are not RSGs, they are of a similar luminosity but with different surface temperature ranging from blue to yellow supergiants.

Besides Type IIP SNe, the only other SN type for which there are a large number of pre-explosion images available are the hydrogen deficient Type Ibc SNe. There have been many attempts to directly identify the progenitors of Ibc SNe in these pre-discovery images, but no success see Van Dyk, Li & Filippenko (2003c); Maund, Smartt & Schweizer (2005); Gal-Yam et al. (2005); Maund & Smartt (2005b); Crockett et al. (2008, 2007). A luminous outburst or eruption from a progenitor star for the type Ibn SN2006jc was detected (Pastorello et al. 2007a; Foley et al. 2007), but the progenitor was not detected in a quiescent phase. The progenitors of Type Ibc SNe could potentially be the classical, massive Wolf-Rayet stars which are initially massive ($\mathrm{M_{ZAMS}} \gtrsim$25-30 $\mathrm{M_{\odot}}$) but then lose their hydrogen envelopes through radiatively driven stellar winds (Crowther 2007). Such mass-loss in massive stars can also be enhanced in massive binary systems.

Alternatively, it has been suggested that progenitors of Type Ibc SNe could be helium stars with ZAMS masses that are too low to allow a *single* star to enter the WR stage. In these cases, the star could have an initial mass of significantly less than $\lesssim 25\ \mathrm{M_{\odot}}$. The lower initial mass limit for a SN progenitor in such a system is uncertain, but is probably around the limit for single stars to produce Fe-cores, or OMgNe cores, around 8$\mathrm{M_{\odot}}$ (Smartt 2009). Such a star could lose its hydrogen envelope through interaction with a binary companion, losing mass by Roche-Lobe overflow or common-envelope evolution. (e.g. Podsiadlowski et al. 1992; Nomoto, Iwamoto & Suzuki 1995; Vanbeveren, Van Bever & Belkus 2007). While theoretically such stars are likely progenitors of Type Ibc SNe, there are no direct hydrogen-free analogues in our own galaxy, which has led to some scepticism as to their existence. The closest such systems we are aware of are V Sagittae systems, WR7a and HD45166, which retain some hydrogen on their surfaces (e.g. Steiner & Diaz 1998; Oliveira et al. 2003; Steiner & Oliveira 2005; Groh, Oliveira & Steiner 2008). However, recently refined measurement of the relative rates of the SN types, together with modelling of the evolution of binary systems, has added weight to the argument that these binary stars represent a significant fraction of Type Ibc progenitors (e.g. Eldridge, Izzard & Tout 2008; Yoon, Woosley & Langer 2010; Smith et al. 2011; Eldridge, Langer & Tout 2011).

In this paper we combine the observed upper limits on the luminosity of individual Type Ibc progenitors with theoretical modelling of a binary population to constrain the progenitor population as a whole. In particular, we aim to determine whether the observed magnitude limits for Type Ibc SN progenitors (and their relative rates) can be reconciled with the observed population of WR stars in the Milky Way and Magellanic Clouds, or whether they are better matched by a model population including binary and single stars.

2 THE LOCAL SN RATE

We have followed the methodology of Paper I in sample selection. The list of supernovae maintained by the IAU Central Bureau for Astronomical Telegrams[2] was searched for all supernovae with a named host galaxy discovered in the two year period from 1 Jan 1998 to 30 March 2012. This sample was then cross matched against the the HyperLEDA galaxy database[3] to identify all supernovae for which the host galaxy had a recessional velocity (corrected for Local Group infall on Virgo) of 2000 km s^{-1} or less, corresponding to a distance limit of 28 Mpc for H_0=72 km $s^{-1}$$Mpc^{-1}$. For SNe where there was not a named host in the IAU catalogue, HyperLEDA was searched for any galaxy within $1.5'$of the SN position. Finally, the coordinates of any SNe which did not have a named host or a galaxy within $1.5'$, and was brighter than magnitude 16 were queried via the NASA Extragalactic Database (NED)[4]. We also cross-checked our sample against the catalogue of Lennarz, Altmann & Wiebusch (2012), by searching for all SNe within 35 Mpc, and manually checking LEDA for the distance to any which were not in our sample. We did not identify any SNe within our distance limit which we had missed, although we note that several SNe (e.g.. SNe 2001fu, 2004ea and 2006mq) which were within our distance limit do not have an associated distance in the Lennarz et al. catalogue.

---

[1] The types Ib and Ic are often difficult to distinguish and their classification is dependent on the phase of the spectra and signal-to-noise. Hence we will often use the common term of type Ibc to cover them jointly.

From this, we obtained a sample of 203 transients over the 14.25 year time period within a 28 Mpc volume, as listed in Appendix A. For comparison, in Paper I we found 141 transients within the same volume over the 10.5 years from 1 Jan 1998 to 30 June 2008, which appears broadly consistent with the number found here : 13.4 SNe yr^{-1} in Paper I, compared to 14.3 SNe yr^{-1} over the extended 14.25 yr period that we consider here. Most of the SNe in the sample have a spectral typing, which are listed in Appendix A together with the recessional velocity of the host. Those SNe which are of uncertain type, or which are otherwise noteworthy, and which were not discussed in Smartt et al. (2009), are discussed individually below.

(i) SN 2010dn

The nature of this transient is still debated. Vinkò et al. (2010) claimed it is an eruptive outburst from a dust-enshrouded LBV, but the spectra bear resemblance to the SN2008S-like transients that are still potentially SNe. (Botticella et al. 2009; Smith et al. 2009; Preito et al. 2010). Berger (2010) set upper limits on the progenitor magnitude of SN 2010dn from archival Spitzer and Hubble Space Telescope (HST) observations of the site of the transient. Smith et al. (2011b) presented a re-analysis of the same HST data, and similarly to Vinko et al. favoured a supernova impostor / LBV eruption as the most likely explanation. Hoffman et al. (2011) report that a MIR detection of 2010dn which is again similar to that found for other SN2008S-like events. We thus leave it in the bin of "Uncertain or non-supernova" in Table A1.

(ii) SN 2009ip

As noted in Table A1, the original discovery of the object we call SN2009ip was an LBV outburst. Smith et al. (2010) and Foley et al. (2011) identified the progenitor as a star of ∼60 $M_\odot$ using archival HST images. A similar transient, UGC 2773 OT2009-1 was discussed by Smith et al., and again was found to have a massive LBV progenitor. These LBV outbursts or eruptions are not included in our rate calculations but are listed, insofar as we are aware of their existence, in Table A1). The star which underwent an LBV eruption to give SN2009ip may now have produced a supernova event, during August-September 2012. If confirmed this is perhaps one of the most interesting supernovae ever to have occurred and is already the focus of intensive study (Mauerhan et al. 2012; Pastorello et al. 2012; Prieto et al. 2012; Levesque et al. 2012). The event occurred outside our timeframe, hence we do not include it as a SN IIn for rate estimates. However, we recognise that SN 2009ip has an extremely important role in linking the progenitor star, SN explosions and the explosion mechanism.

(iii) SN 2008iz

The SN was discovered at radio wavelengths (Brunthaler et al. 2009), and has been the subject of extensive radio follow-up (Brunthaler et al. 2009; Marchili et al. 2010). Fraser et al. (2009) observed the SN with Gemini+NIRI, and initially claimed that the SN could not be seen in the NIR. However a careful reanalysis by Mattila et al. (2012a) shows that the SN can be seen when image subtraction techniques are used. They find a likely very high line of sight extinction, but probably less than $A_V \approx 10$. As this was a SN which was not, and could not, have been discovered by current optical surveys we do not consider it for our rate estimates. Its type is also currently unknown which would not enhance our measurements. It does serve as a reminder on the number of SNe that may be occurring in high star-formation rate locations which are extinguished (Mattila et al. 2012a,b; Botticella et al. 2012).

(iv) SNe 2008ge, 2008ha, 2010ae & 2010el.

The SNe have all been identified as SN 2002cx-like. 2008ha is perhaps the most unusual SN considered in the sample; it showed low velocities and a faint absolute magnitude, and its interpretation is still contentious. Valenti et al. (2009) suggested that the SN was the faint core-collapse of a massive progenitor on the basis of its spectral similarity to SN 2005cs, however Foley et al. (2009) and Foley et al. (2010a) have questioned this interpretation and proposed that it was in fact a thermonuclear SN. Foley et al. argue that intermediate mass elements, whose absence Valenti et al. claimed was counterfactual to a Type Ia scenario, are in fact present. Valenti et al. noted that SN 2008ha appears to be related to the class of SN 2002cx-like SNe (Li et al. 2003), and suggested that all of these may be related to Type Ibc SNe, possibly with fallback of the inner most ejecta onto the compact stellar remnant. While a fallback explosion model has been constructed for SN 2008ha (Moriya et al. 2010), it requires a progenitor which has a mass lower than that typically thought to undergo this process (Heger et al. 2003). Remarkably, three other SNe besides SN 2008ha in this sample have been suggested to be SN 2002cx-like. The discovery of this many SN 2002cx-like SNe in such a short period could suggest that a significant number of these events have gone undetected previously, likely due to their faint magnitude. For one of them, SN 2008ge, Foley et al. (2010b) have analysed pre-explosion data, and find no evidence of either ongoing star formation, or the young massive stellar population which would be expected to be found in the environment of a core-collapse SN. While the nature of these

---

[2] http://www.cbat.eps.harvard.edu/lists/Supernovae.html

[3] http://leda.univ-lyon1.fr/

[4] http://ned.ipac.caltech.edu/



Figure 1. Percentage of CCSNe which are of a particular type as found in this work.

SNe is still open to debate, at present the balance of evidence seems to favour a thermonuclear interpretation, and so we have excluded them from the core-collapse sample on this basis. However, given recent evidence that the class of SN 2002cx-like SNe may be significantly more heterogeneous than previously thought (Narayan et al. 2011) and may not all share the same explosion mechanism, a more detailed search for the progenitors of these SNe would be a worthwhile endeavour.

(v) SN 2008eh

As far as we are aware, there is no published spectrum of SN 2008eh. Hence we have followed the lead of Horiuchi et al. (2011) and assumed the SN is a core-collapse event (probably Type Ibc) on the basis of its absolute magnitude and lightcurve, and proximity to a HII region in a spiral galaxy. Nonetheless, we have excluded this SN from the rate calculations.

We summarise the relative rates of different core-collapse SNe within our distance limit in Table A1. Discarding all SNe which were designated as Type Ia (55 SNe, or 30.2 per cent of the total classified sample) along with SN impostors and unclassified transients, we find relative rates for the various types of core-collapse SNe as listed in Table 1 and shown in Figure 1. SNe which were classified as a "Type Ibc" were divided between the Type Ib and Ic bins, according to the observed ratio of these subtypes (9:17). Type II SNe were distributed between the IIP, IIL, IIb and IIp-pec bins in the ratio 55:3:12:1. Uncertainties were calculated from the standard error on a Poissionian distribution (i.e. as $\sqrt{N}$). We have made no attempt to further subdivide the Type IIb SNe into the "compact" and "extended" subtypes proposed by Chevalier & Soderberg (2010).

Our relative rates compare favourably with those found in Paper I, the most noticeable difference being the increase in Type IIb SNe at the expense of Type Ibc events. A possible explanation for this is that SNe are now discovered (on average) sooner after explosion, and so Type IIb SNe which previously would have been detected late and classified as a Type Ibc are now correctly identified.

Smith et al. (2011) measured SN rates from the LOSS sample, based on those SNe selected for the luminosity function analysis by Li et al. (2011), with a total of 106 CCSNe within 60 Mpc. This is smaller than we present here, but is a much more homogeneous sample and is a well controlled SN survey. We emphasise that we are assimilating all discoveries and not carrying out a controlled SN search, but the results should be compared given the large numbers available.

Smith et al. did not separate out the IIpec category, but did separate the Ibc-pec SNe. To facilitate a comparison between our rates, we re-distributed the Type Ibc-pec SNe from Smith et al. between the Ib and Ic categories (weighted by the Type Ib:Ic ratio), and the Type IIpec SNe from this work between the Type IIP, IIL and IIb categories. We find a good agreement between the Smith rates and those presented in this paper for the Type IIb, Ib and Ic SNe. We find a Type IIP rate which is 7.3 per cent higher than that derived by Smith et al, although the uncertainties of Smith et al. and this work do overlap. If we combine the Type IIL and IIP rates, then the discrepancy is smaller, raising the possibility that some of our Type IIP SNe may in fact be Type IILs (and vice-versa for Smith et al.). Smith et al. suggest that the LOSS rates are more reliable due to the SNe all coming from the same survey with well controlled and known selection effects and good spectroscopic and photometric coverage. The latter is certainly important to distinguish between II-P and II-L SNe. In Paper I we attempted to classify II-P SNe from literature, archive and web data searches and classified any type II SN that had a constant magnitude phase for more than 30 days a type II-P. This of course is not as homogeneous a dataset as that from LOSS. There are 23 SNe in common between the Smith et al. sample and that presented in Paper I (there are no extra SNe to add from 2008-2012, as the LOSS sample stops in 2006). Of those, 19 are identical classifications and two have type II classifications by us and II-P classifications by LOSS. Including those two updated LOSS classifications here would make the II-P discrepancy worse. The other two were fairly trivial differences : Ic vs Ibc/IIb (2004ao) and Ibc vs Ib (2004qn) in Paper I and LOSS respectively. Hence there is no evidence we can see that points to systematic miss-classifications in the smaller volume sample presented in Paper I and supplemented here.

The only SN type for which the rates and their uncertainties do not overlap at all are the Type IIn SNe, where Smith et al. find a higher rate than we do ($8.8^{+3.3}_{-2.9}$ per cent versus 2.4 ± 1.4 per cent). Smith et al. attribute this to the fact that LOSS has better and more extensive spectroscopic and photometric coverage. This is difficult to assess as the global community certainly has extensive resources to bear on SNe within 28 Mpc. Classification results are rapidly reported, data are published on reasonable timescales, and we have made attempts to search archives and request private information where we can. Hence it's not clear to us that this statement in Smith et al. is necessarily correct. The Li et al. sample was produced for luminosity function work, which requires SNe to be "in season" and full lightcurve coverage from before explosion. This is essential for luminosity function measurements, but not clearly correct for rate estimates. For example, long duration plateau SNe may be preferentially excluded as their "season" lasts longer than other types. It's still possible that the high IIn rate in LOSS is due to them being brighter and being preferentially discovered between 30-60 Mpc compared with < 28 Mpc. However overall, the agreement is reasonable despite the differences in the construction of the two samples.

As described in Paper I for all SNe, we searched the

Table 1. The relative frequency of core-collapse SN types discovered between 1998-2012.25 (14.25 yrs) in galaxies with recessional velocities less than 2000 kms^{-1}, as listed in Table A1. The values in parentheses in the Number column include SNe of indeterminate subtype as discussed in the text.

SN Type	Number	Relative rate (per cent)	LOSS (per cent)
IIP	55 (70.5)	55.5 ± 6.6	$48.2\,^{+5.7}_{-5.6}$
IIL	3 (3.8)	3.0 ± 1.5	$6.4\,^{+2.9}_{-2.5}$
IIn	3 (3)	2.4 ± 1.4	$8.8\,^{+3.3}_{-2.9}$
IIb	12 (15.4)	12.1 ± 3.0	$10.6\,^{+3.6}_{-3.1}$
IIpec (87A-like)	1 (1.3)	1.0 ± 0.9	...
Ib	9 (11.4)	9.0 ± 2.7	$8.4\,^{+3.1}_{-2.6}$
Ic	17 (21.6)	17.0 ± 3.7	$17.6\,^{+4.2}_{-3.8}$
Total	100 (127)		

HST archives using the Querator[5] and MAST[6] webpages. For core-collapse SNe we find that out of 127 SNe, 41 of them had HST images taken before discovery (i.e. 32 per cent) with the SN falling on the HST camera field of view. For type Ia SNe the fraction was 20 out of 55 (i.e. 36 per cent). Many more host galaxies of SNe had observations taken before explosion, but the SNe fell outside the small fields of view of the HST cameras (16 per cent of core-collapse and 16 per cent of type Ia ; see Table A1). We also searched the archives for the ESO telescopes[7], Gemini North and South telescopes[8], the Subaru Telescope[9], the William Herschel Telescope (WHT)[10] and the Canada-France-Hawaii Telescope (CFHT)[11] for any SNe in a host with a recessional velocity of <1000 km s^{-1}. There are other 8 meter class telescopes which could be of use for progenitor searches, such as the Keck telescopes and the Large Binocular Telescope, however the absence of publicly accessible data archives precludes this.

3 ESTIMATING PROGENITOR MASSES FROM SN RATES

Smith et al. (2011) used their observed SN rates to estimate progenitor mass ranges, given a standard IMF and assuming that more massive stars will undergo greater mass loss. They reached the conclusion that a simple single star initial mass prescription is unable to reproduce the observed SN rates, and that binary evolution is a crucial factor for Type Ibc SNe. This is based on the relatively high rates of Ibc SNe and the fact that the mass range for classical, massive WR stars cannot reproduce the rates from Salpeter IMF arguments (also see discussion in Smartt 2009). This problem was originally noted by Podsiadlowski et al. (1992) and Nomoto, Iwamoto, & Suzuki (1995) who proposed that around 30 per cent of stars above 8M$_\odot$ could lose mass in interacting binaries and produce the Ibc SN population. Here we use a more detailed stellar population model to reproduce the observed rates. The main difficulty is attaching an observed SN type to the end point of the stellar model and the physical mass of H and He left in the model stars.

To reproduce the SN rates with a binary population, we use synthetic stellar populations from the Binary Population and Spectral Synthesis code (BPASS). This code is able to create stellar populations of both single and binary stars, and is described in detail in Eldridge, Izzard & Tout (2008), Eldridge & Stanway (2009) and Eldridge, Langer & Tout (2011). In these papers it is shown that populations including binaries are better able to reproduce observed stellar populations, both resolved and unresolved.

We have used the BPASS code to predict the relative rates of different types of SNe determined in Section 2 and listed in Table 1. Our fiducial progenitor populations are a solely single-star population, and a mixed binary/single-star population with one single star for every binary. The models are computed for two metallicities, Z = 0.008 and 0.020, corresponding to the Large Magallenic Cloud abundance and the Solar abundance respectively. We average over these values as the SNe in our sample occur in nearby spiral galaxies which will have metallicities between these two extremes (see discussion in Paper I). We also note that at lower metallicities quasi-homogeneous evolution may become important (Yoon, Langer & Norman 2006; Eldridge, Langer & Tout 2011) but we do not currently include this pathway. We consider all stars from 5M$_\odot$ to 120M$_\odot$ in both single and binary populations. For the binary populations we include a range of mass ratios for the secondary star (q = 0.3, 0.5, 0.7 and 0.9) and a range of initial separations from $\log(a/R_\odot) = 1$ to 4. According to our models, approximately two-thirds of these binaries interact. This is consistent with the recent observations of Sana et al. (2012), although those are measurements of O-stars with ZAMS masses of 20M$_\odot$ and greater.

The precise configuration of a star at the moment of core-collapse, and how this determines the resulting SN type is still quite uncertain (e.g. Heger et al. 2003; Eldridge & Tout 2004b). Stellar evolution codes calculate model stars with quantitative residual envelope masses of H and He and are given SN types by assuming values of the minimum masses required to give SNe of types II and Ib respectively (for example, see the discussion in Dessart et al. 2011). Hence we look at this problem in a different way, we insist that the model populations must reproduce the observed relative SN rates in Table 1, as in Eldridge, Langer & Tout (2011). Each star in the ensemble is evolved to the end of core carbon burning. At this point, a resulting SN type is assigned to the model, based on the mass of the hydrogen envelope, the H to He mass ratio, and the fraction of He in the ejecta. Then we can determine what the minimum mass of H is to produce the total numbers of type II and type II-P populations respectively. As suggested by Smith et al. (2011), we find that a single star population cannot produce the observed rates of Ibc SNe

---

[5] http://archive.eso.org/querator/
[6] http://archive.stsci.edu/
[7] http://archive.eso.org/cms/
[8] http://cadcwww.dao.nrc.ca/gsa/
[9] http://smoka.nao.ac.jp/
[10] http://casu.ast.cam.ac.uk/casuadc/archives/ingarch
[11] http://cadcwww.dao.nrc.ca/cfht/

A mixed population of binary and single stars (with one single star for every binary) produces end stellar points that can match the relative rates of type Ibc vs. II SNe *if* we assume that the final hydrogen and helium masses given in Table 2 are what is required to produce the various types. We find that a Type II SN requires at least $0.003\mathrm{M}_{\odot}$ of H in the ejecta, while a Type IIP SN must have a H to He mass ratio of greater than 1.05 and that the total mass of hydrogen is greater than $1.5\ \mathrm{M}_{\odot}$. While this mass of $0.003\mathrm{M}_{\odot}$ of H is at first sight low, it would be the minimum for a IIb SNe. The synthetic spectra calculations of Dessart et al. (2011) show that even $0.001\mathrm{M}_{\odot}$ can produce visible hydrogen lines (Hα in particular), as long as the surface mass fraction is higher than 0.01.

For a Type Ib SN we require the mass fraction of helium in the ejecta to be greater than 0.61, while for a Type Ic the He fraction must be less than this. Comparing these mass values to those in Eldridge, Langer & Tout (2011) - which were based on the rates of Paper I - we find that they have changed by a similar percentage as the SN rates between Paper I and this paper. The most uncertain value is the amount of hydrogen that can be hidden in a type Ibc SNe, the value here being lower by a factor of 17. This difference is chiefly due to the larger fraction of Type IIb SNe in this work, which causes the minimum required H mass for a Type II SN to be lower.

We appreciate that this fairly simple and well defined classifications of the progenitor models are, in reality, complicated by two factors. Branch et al. (2002) show that some Ib SNe may well contain some residual hydrogen, given good quality spectra and model photosphere fits. Dessart et al. (2012) discuss that it may be quite possible to get a Ic SN with helium in the progenitor star, when He I is not excited due to weak mixing of ^{56}Ni into the He rich regions.

In summary we can reproduce the observed rates in Section 2, with a mixed population of single stars and binaries with one single star for every binary system, as long as we attach the resulting progenitor stars into SN classification bins with the envelope mass fractions as reported in Table 2. This means that one third of all progenitors have had their envelopes stripped by binary interactions via either Roche lobe overflow or common envelope evolution (consistent with results found in Smith et al. 2011).

Finally it has been suggested that SNe that form black holes directly do not produce any SN display. Heger et al. (2003) estimate that there will be no optical display if the final helium core mass is greater than $15\mathrm{M}_{\odot}$; in our models this corresponds to a remnant mass of $8\mathrm{M}_{\odot}$. We have created a synthetic population in which we remove all core-collapse events with a remnant mass $> 8\mathrm{M}_{\odot}$ (approximately 10 per cent of all SNe). Even without these events, we find that the same parameters in Table 2 can be used to reproduce the observed SN rates, so long as we increase the fraction of binaries in the population.

Table 2. The required parameters for a star to give rise to a specific SN type assuming all SN are observable.

SN Type	Final Mass/$M_{\odot}$	M_{COcore} /$M_{\odot}$	M(H) /M(He)	M(H) /$M_{\odot}$	M(He) /M_{ejecta}
IIP	> 2	> 1.38	$\geqslant 1.05$	> 1.5	-
II	> 2	> 1.38	< 1.05	> 0.003	-
Ib	> 2	> 1.38	–	$\leqslant 0.003$	$\geqslant 0.61$
Ic	> 2	> 1.38	–	$\leqslant 0.003$	< 0.61

4 LIMITS ON TYPE IBC SN PROGENITORS FROM ARCHIVAL IMAGING

Alongside estimates from the relative rates of CCSN types, direct detections and upper limits on progenitors from archival data can help constrain their progenitor population. In the following section we summarise the published limits for Type Ibc progenitors within a recessional velocity limit of 2000 km s^{-1}, and extend this sample with limits on an additional six SNe.

The general technique used is the same as that described in Smartt et al. (2009). We align a pre-explosion image to a post-explosion image by means of background stars in the vicinity of the SN. We then search for a source coincident with the SN in the pre-explosion image. If a coincident source is identified, then its magnitude can be measured, otherwise (as in all of the following cases) either pixel statistics or artificial star tests are used to set a limiting magnitude for a non-detection. All magnitudes quoted here are Vegamags.

4.1 2000ew

SN 2000ew was discovered by amateur astronomers (Puckett, Langoussis & Garradd 2000). The SN was initially spectroscopically classified as a Type Ia SN (Dennefeld & Patris 2000), before Filippenko, Foley & Modjaz (2000) obtained a second spectrum which showed the SN to be of Type Ic. A Tully-Fisher distance of 18.2 Mpc has been measured for the host galaxy of SN 2000ew, NGC 3810. As the coordinates of NGC 3810 are on the outskirts of the Virgo cluster (11h41m, +12°28′) the kinematic distance estimate (13.4 Mpc) may not be reliable. We hence adopt the Tully-Fisher distance as a more conservative estimate.

Maund & Smartt (2005b) set a limit of F606W>24.6 on the progenitor of SN 2000ew from pre-explosion WFPC2 imaging, while Van Dyk, Li & Filippenko (2003a) find a similar limit of F606W>24.7. Maund & Smartt estimate the reddening towards the SN to be E(B-V)=0.01, but as this is less than the foreground value from NED (E(B-V)=0.044) we adopt the latter. The F606W absolute magnitude is listed in Table 4.

Late time NIR spectra of SN 2000ew were presented by Gerardy et al. (2002), these are can help further elucidate

Table 3. HST Observational Data

Filter	Date	Exposure time	
SN2001ci			
WFPC2	F547M	1999-03-04	320
WFPC2	F606W	2001-01-21	560
WFPC2	F658N	1999-03-04	1200
WFPC2	F658N	1998-11-26	8900
WFPC2	F814W	1998-11-26	800
WFPC2	F814W	1999-03-04	140
SN2003jg			
WFPC2	F450W	2001-08-02	460
WFPC2	F606W	1994-06-21	160
WFPC2	F814W	2001-08-02	460
SN2004gn			
WFPC2	F658N	2011-03-02	1400
WFPC2	F606W		600
SN2005V			
ACS	F658N	2004-04-10	700
	F814W		120
NICMOS	F160W	2002-12-04	96
	F187N		640
	F190N		768
WFPC2	F450W	2001-07-06	460
	F814W		460
	PC F606W	2001-01-05	560

the nature of the progenitor. Carbon monoxide (CO) emission was observed in the spectrum, albeit at a lower velocity (∼2000 km s^{-1}) than might be expected for a progenitor with a 2.1 $M_\odot$C+O core (Iwamoto et al. 1994). While the CO velocity could be interpreted as arising from a lower energy explosion, perhaps pointing to a less massive progenitor, it may also be explained with an asymmetric explosion with a largest component of the velocity perpendicular to the line of sight. We also note that Gerardy et al. (2002) also found narrow HeI and [FeII] lines, but no Brackett series lines, in their spectrum which they attribute to H-poor material shed by the progenitor star prior to core-collapse.

4.2 2001B

SN 2001B was discovered in IC 391 by Xu & Qiu (2001), and initially classified as a Type Ia SN by Matheson et al. (2001) before Chornock & Filippenko (2001) reclassified the SN as a probable Type Ib. A lightcurve was presented by Tsvetkov (2006) which reached a maximum of -17.5 in V, supporting the classification of SN 2001B as a Type Ib. IC 391 has a Tully-Fischer distance of 25.5 Mpc, and a recessional velocity distance of 25.5 Mpc (from NED).

Van Dyk, Li & Filippenko (2003c) claimed the tentative detection of a progenitor candidate for SN 2001B from an alignment to a ground-based natural seeing image. Subsequently Maund & Smartt (2005b) presented late time HST imaging showing that the SN was offset from the candidate. Maund & Smartt set an upper limit of F555W=24.30±0.15 for the progenitor of SN 2001B, which is in agreement with the limit suggested by Van Dyk et al. if their progenitor candidate was *not* the progenitor. Maund & Smartt estimate a reddening of E(B-V)=0.102±0.030 from nearby supergiant stars, which is slightly lower than the line of sight extinction of E(B-V)=0.14 from NED, we hence adopt the latter as the larger value. The absolute magnitudes are reported in Table 4.

4.3 2001ci

SN 2001ci was discovered as part of the Lick Observatory and Tenagra Observatory Supernova Searches (LOTOSS) by Swift (2001). It was noted at discovery that the SN had a low absolute magnitude, and it was originally suggested that the transient may be a SN impostor. Filippenko & Chornock (2001) subsequently obtained a spectrum of the SN, and classified it as a highly reddened Type Ic SN, with a line-of-sight extinction of $A_V \approx$ 5–6 mag.

While NGC 3079 had been observed with HST prior to the SN explosion as detailed in Table 3, the images are not sufficiently deep to have a reasonable expectation of seeing a progenitor given the high extinction. In addition, we were unable to locate any high resolution post-explosion imaging for SN 2001ci. NGC 3079 was observed with HST+WFPC2 in the F300W filter on 2001 Dec 9 (≈6 months after the SN explosion) and the SN coordinates fall in the field of view, but it is not visible (as SNe are typically only bright at UV wavelengths for some weeks after explosion). We searched the CFHT, Gemini, Subaru, ING, TNG and NOAO archives, but no images of SN 2001ci were found. We have hence not considered SN 2001ci any further.

4.4 2002ap

SN 2002ap was discovered in M74 by amateur astronomers (Nakano et al. 2002), and rapidly classified as a type Ic SN (Meikle et al. 2002). SN 2002ap was extremely energetic, and bore a striking spectroscopic similarity to the "hypernovae" 1997ef and 1998bw (Mazzali et al. 2002; Gal-Yam, Ofek & Shemmer 2002). The explosion is thought to be moderately asymmetric, both from early time spectropolarimetry (Kawabata et al. 2002; Wang et al. 2003) and modelling of the late time nebular spectra (Mazzali et al. 2007).

The first attempt to detect the progenitor of SN 2002ap was made by Smartt et al. (2002), who did not find a progenitor in ground-based pre-explosion imaging. A subsequent study by Crockett et al. (2007) used even deeper pre-explosion images, which still did not yield a progenitor detection. The limits for SN 2002ap remain the deepest limits for a hydrogen-deficient SN to date. The 5σ limiting magnitudes found by Crockett et al. are B= 26.0±0.2 mag, and R=24.9±0.2 mag. The limits from Smartt et al. (2002) are much shallower, with the exception of their U-band image, which is of interest for constraining a hot progenitor, and had a 3σ limit of 21.5 mag. Hendry et al. (2005) considered the various measurements of the distance towards the host galaxy for SN 2002ap, M74, and adopted an average value of 9.3 ± 1.8 Mpc, which we also adopt (as in Crockett et al. 2007). The total extinction (Milky Way and host galaxy) towards SN 2002ap is given by Takada-Hidai, Aoki & Zhao (2002) as E(B-V)=0.09±0.01, from a high-resolution spectrum of the NaID lines.

4.5 2003jg

SN 2003ig was discovered by LOTOSS (Graham & Li 2003), and classified by Filippenko & Chornock (2003) as a Type Ic SN approximately a week after maximum light. The SN exploded in NGC 2997, which has a Tully-Fisher distance of 12.2 ±0.9 Mpc (Hess et al. 2009), which is in good agreement with the recessional velocity based distance of 12.7 Mpc. The foreground extinction towards NGC 2997 from Schlegel, Finkbeiner & Davis (1998) is E(B-V)=0.109 mag.

Post explosion images of SN 2003jg were obtained in several filters with HST+ACS/HRC on 2003 Nov 18. Pre-explosion WFPC2 images of NGC 2997 as detailed in Table 3 were used to search for the progenitor. The images in each filter consisted of a cr-split pair, these were combined with the CRREJ task in IRAF to reject cosmic rays. The pre-explosion WFPC2 F450W and F814W images were taken at the same pointing, with the SN position falling on the WF3 chip. To accurately identify the SN position in these, the F814W image was aligned to the post explosion ACS/HRC F814W image. The SN position in pre-explosion F606W (on the PC chip) image was determined by aligning it to the post-explosion ACS/HRC F555W image.

20 sources were identified common to both the F814W ACS and F814W WFPC2 images, and used to derive a general transformation with an rms error of 17 mas. The SN position was measured in the ACS F814W image (j8qg10021_drz) to be (704.530, 259.953), with an uncertainty of 0.3 mas, which was transformed to the pixel coordinates of the WFPC2 F814W image (u6ea3803r_c0f), where it was found to be 246.10, 164.15, with an uncertainty of 0.2 pixels

The pre- and post-explosion F606W WFPC and F555W ACS images were aligned with an rms error of 16 mas from a general transformation to 29 common sources; the SN position was measured in the post explosion image to lie at pixel coordinates 704.844, 259.740 in j8qg10050_drz with an rms error of 1 mas. The SN position was transformed to the pre-explosion WFPC2 image, where it was found to have pixel coordinates on the PC chip of 334.99, 166.94 in u29r1301t_c0f, with a total uncertainty of 0.3 pixels.

Inspection of the transformed SN position in the pre-explosion frames did not reveal any coincident source, as can be seen in Fig. 2. HSTPHOT was run with a 3σ detection threshold on the pre-explosion images, but did not detect a source coincident to the SN at this significance. We calculate 5σ limiting magnitudes for the progenitor of SN 2003jg to be F450W>24.63, F606W>24.63 and F814W>23.70 using the technique described in Crockett et al. (2011).

4.6 2004gk

SN 2004gk was discovered by Quimby et al. (2004) and classified as a Type Ic SN near maximum light. The host galaxy of SN 2004gk, IC 3311, is at an inclination angle of 90deg (from LEDA), rendering any search for the progenitor optimistic at best. We note however that Elmhamdi et al. (2011) report a host galaxy extinction of A_V=0.372 mag from low resolution spectroscopy, while from photometry the SN does not appear to be heavily reddened. It is hence plausible that while IC 3311 is edge on to our line of sight, SN 2004gk may have exploded relatively close to the edge of the galaxy. The galaxy has a negative recessional velocity as it lies within the Virgo supercluster, however the Tully-Fischer distance to the galaxy from LEDA is ∼20 Mpc. Unfortunately, there are no pre-explosion HST images of the galaxy, while natural-seeing ground based imaging is not useful at such distances. We have hence not considered SN 2004gk any further in the following.

4.7 2004gn

SN 2004gn was discovered with the Katzman Automatic Imaging Telescope (KAIT) as part of the Lick Observatory Supernova Search (LOSS) close to the nucleus of NGC 4527 (Li, Yamaoka & Itagaki 2004). The spectral classification of SN2004gn was not reported in an IAU CBET. Private communication from the KAIT team (B. Cenko) confirms that it is a H-deficient SN and a Ibc classification is reasonable. [12]. A Nearby Supernova Factory spectrum [13] shows it to be a Ibc similar to SN1990B at 90 days, which is a Ib. As the SN was discovered late, we cannot rule out that there was residual hydrogen at early times that has disappeared, but with the lack of early data we assume a classification of Ibc. We found very late time spectroscopy of the SN location from the WHT in the ING archive, however no flux from the SN was present and so this was of no use for this work.

Besides SN 2004gn, NGC 4527 was also host to the peculiar Type Ia SN 1991T. Subsequently, the galaxy has been the subject of numerous Cepheid studies with HST to obtain an independent measure of the distance. The most recent of these (Saha et al. 2006) found a distance of 14.2±1.3 Mpc towards NGC 4527. SN 1991T itself implies a distance of 14.1 Mpc from fits to the multi-colour light curve (Jha, Riess & Kirshner 2007). We have adopted the average of these two values (14.2±1.3 Mpc) as the distance towards the host. The extinction towards SN 2004gn is poorly constrained, as we have no spectra, or information on phase to allow us to compare the magnitude to other Type Ibc SNe. Hence we adopt the foreground extinction from NED, A_V=0.07 mag, as a lower limit to the extinction.

Broad-band pre-explosion images for SN 2004gn (as listed in Table 3) consist of two 300s WFPC2 F606W images, where the SN falls on the WF3 chip. Unfortunately no post explosion images of the SN were discovered during our archival search. As an alternative to a direct alignment between pre- and post- explosion frames, we attempted to use three isolated SDSS point sources to register the combined WFPC2 WF3 image to the celestial WCS. Attempting to solve this with only three sources means we are over fitting the data, and it does not make sense to quote an rms error in the alignment. However, by inspection of the positions of other SDSS sources (which were not used in determining the transformation, as they are extended), we believe that the accuracy of the registration is $\lesssim 1''$.

The RA and Dec of SN 2004gn as reported on the IAU webpages correspond to pixel coordinates of 77.8, 204.0 on the WF3 chip (using the new WCS solution). We ran HSTPHOT with a detection threshold of 3σ. Within a $1''$radius of the reported SN coordinates (shown in Fig. 3), we find a

---

[12] http://astro.berkeley.edu/bait/public_html/2004/sn2004gn.html

[13] http://www.rochesterastronomy.org/sn2004/sn2004gn.jpg



(a) Pre-explosion WFPC2/WF3 F450W image (b) Pre-explosion WFPC2/PC F606W image (c) Pre-explosion WFPC2/WF3 F814W image (d) Post-explosion ACS/HRC F814W image

Figure 2. Pre-explosion HST+WFPC2 images of the site of SN 2003jg, plus a post-explosion HST+ACS image. Scale and orientation for each panel is identical to that indicated in the post-explosion image. The circle in the upper right corner of each panel has a radius equal to 5 times the rms error in the position of the supernova in that image.



Figure 3. Pre-explosion HST WFPC2 image of the site of SN 2004gn, taken with the F606W filter. As no post-explosion images were available, we show a 1″(dashed line) and 2″(solid line) radius around the published coordinates for the SN.

single source detected above the 5σ level, at a magnitude of F606W=25.4. Six other sources are detected above 3σ, but below 5σ, and with magnitudes between 25.4<F606W<25.9. Within the larger annulus, seven sources are detected above the 5σ level, the brightest of which has a magnitude of F606W=24.2. The brightest source has a large χ value, and a sharpness parameter which is more negative than would be expected for a single star. In fact, all of the sources detected by HSTPHOT have negative sharpness, suggesting that these are not stellar, but rather that HSTPHOT is detecting unresolved background structure. Nevertheless, we adopt the magnitude of the brightest source detected by HSTPHOT as our limit for the progenitor magnitude, i.e. F606W>25.4. The F658N images were examined, but no Hα excess was seen at the SN position.

4.8 2004gt

Monard (2004) discovered SN 2004gt in NGC 4038; the SN was spectroscopically classified as a Type Ic SN. Kinugasa, Kawakita et al. Yamaoka (2004) suggested that it was difficult to distinguish between the Ib and Ic subclasses therefore we take the conservative approach and label the SNe as Type Ibc. NGC 4038 is part of a well the studied interacting galaxy pair Arp 244, better known as the Antennae galaxies. Despite extensive observations of the Antennae, their distance is still a matter of some debate in the literature. Saviane, Hibbard & Rich (2004) derived a distance of 13.8±1.7 Mpc from the tip of the red giant branch (TRGB) using HST WFPC2 imaging. This value was revised by Saviane et al. (2008) to 13.3±1.0 Mpc, based on deeper images from HST+ACS. The TRGB distance is considerably closer than that expected given the recessional velocity of Arp 244, which implies a distance of ∼20 Mpc. The discovery of the Type Ia SN 2007sr in NGC 4038 provided an independent measurement of the distance from a fit to the SN lightcurve, yielding a value of 22.3±2.8 Mpc, in disagreement with the TRGB estimate (Schweizer et al. 2008). Schweizer et al. suggest that Saviane et al. have misidentified the TRGB, and present a re-analysis of the HST data used by the latter which brings the TRGB distance into accord with that from SN 2007sr. We follow the suggestion of Schweizer et al. in adopting 22±3 Mpc as the distance for NGC 4038, based on an average of the values from the recessional velocity, re-calibrated TRGB, and SN 2007sr. The extinction towards SN 2004gt has been taken as E(B-V)=0.07±0.01, following the lead of Maund, Smartt & Schweizer (2005).

Both Maund, Smartt & Schweizer (2005) and Gal-Yam et al. (2005) presented limits on the progenitor of SN 2004gt from pre-explosion HST images. Maund et al. found limits of F336W > 23.04, F439W > 24.56, F555W > 25.86 and F814W > 24.43. The limits found by Gal-Yam et al. are similar, but as their values are quoted in $UBVI$ rather that the HST flight system these cannot be compared directly. Gal-Yam et al. also present a limit from a pre-explosion STIS image of >23.4 for the far-UV. Using this new distance and extinction we determine upper limits for the absolute magnitudes for the progenitor of SN2004gt in Table 4.

4.9 2005V

The host galaxy of SN 2005V, NGC 2146, is a starburst galaxy, and is thought to have a large population of recently formed massive stars. However, the absence of an overabundance of supernova remnants may indicate that the starburst is at an earlier stage than similar galaxies such as M82 (Tarchi et al. 2000). The kinematic and Tully-Fischer estimates for the distance towards NGC 2146 agree well (17.2 and 16.9 Mpc respectively), however the sosies estimate is significantly larger at 27.7 Mpc. As NGC 2146 appears to have a slightly irregular morphology, it is possible that the sosies method is not a reliable measure in this case. We have hence discarded this measure, and adopted the mean of the kinematic and Tully-Fischer distances.

Mattila et al. (2005) discovered SN 2005V in the near infrared, although the early time J-H and H-K colours of the SN do not support very high levels of extinction. Mattila et al. find J-H=0.1 mag, H-K =0.2 mag for SN 2005V, which is indistinguishable from the the typical J-H and H-K colours of Ibc SNe (Hunter et al. 2009). Taubenberger et al. (2005a) classified the SN as a Type Ibc, but with a strong NaID absorption (EW = 5.5 Å) and a red continuum which does indicate significant reddening. Unfortunately there is no published photometry of SN 2005V, so we cannot compare the optical colours and absolute magnitude to other Type Ibc SNe. Using the relation of Turatto et al. (2003) between NaD absorption and reddening, we infer E(B-V)$\sim$0.9, however we are somewhat sceptical of the veracity of this estimate due to the large scatter in the relation (Poznanski et al. 2011). Nonetheless we adopt E(B-V) as the conservative option, as an *over*estimate of A_V will only serve to make our limits *less* restrictive. On a more qualitative level, we also note that the SN is close to the nucleus of the galaxy, that a large number of dust lanes are seen in the HST images, and that the SN position is offset by $\sim$0.15 $''$from one of these lanes.

Pre-explosion WFPC2 images as listed in Table 3 were downloaded from the HST archive. Cr-split pairs were combined with CRREJ to remove cosmic rays. The SN position lies on the PC chip in the F606W image, and on the WF3 chip in the F450W and F814W filter images, as shown in Fig. 4. To identity the SN position in the pre-explosion images, we used images of the SN obtained with the ACS/HRC on 2005-08-08.

The F606W pre-explosion was aligned to the F555W post-explosion image. 26 sources were used to fit a general transformation with an rms error of 0.164 WFPC2/PC pixels, corresponding to 8 mas. The SN position was measured in j90n04020_drz to be 578.770,609.971 with an error of $\ll$1 mas. The SN position was transformed to the pre-explosion image, where it was found to have pixel coordinates 331.27,411.27 in u67n1801r_c0f.fits, with an uncertainty of 0.16 pixels. HSTPHOT did not detect a coincident source at the 3σ level, and we calculated the limiting 5σ magnitude as before to be F606W>24.76.

The F814W pre-explosion image was aligned to the post-explosion F814W image with 17 sources, for an rms error of 0.17 WFPC2/WF pixels, or 17 mas. The SN position was measured in the ACS/HRC F814W image to an accuracy of 1.5 mas, and transformed to the pre-explosion WFPC2 F814W image. We derived transformed coordinates of 140.19, 94.01 for the SN in u6ea2803r_c0f. No source was evident at the SN location, and HSTPHOT detected no source at the 3σ level or above in either the F450W or F814W images. We calculate 5σ limiting magnitudes for the progenitor of SN 2005V of F814W>22.80, F450W>24.05.

The ACS pre-explosion images were also examined. The F814W image showed no source coincident with the SN, and as the limit is comparable to that from the WFPC2 F814W image, this was not considered any further. The F658N image (narrow-band Hα) showed flux at the SN position, but as there was no corresponding continuum observations, it was unclear whether this flux comes from Hα emission or was simply from the high background levels. The NICMOS data were taken with the NIC3 camera, which has a pixel scale of 0.2$''$. The narrow band images were not examined in details, while the broad band F160W image was too shallow to be of use, especially for constraining a compact (and presumably hot) progenitor.

4.10 2005at

SN 2005at exploded in NGC 6744, at a distance of 14.1$\pm$2.8 Mpc. Unfortunately the SN suffers from a high level of extinction ($A_V \sim 2.3 \pm 0.3$, E. Kankare, private communication), while the pre-explosion data consists of ground-based imaging (from VLT+FORS,VIMOS,ISAAC and the ESO 2.2m+WFI) rather than from HST. Using the ESO exposure time calculators[14] we estimated 5σ limiting magnitudes for the pre-explosion images, and took these together with the distance modulus and extinction $A_V \sim 2.3$ to estimate the absolute magnitude reached be the pre-explosion data. As the absolute magnitude reached is between -9 and -11 mag, which is between 1.5 and 3.5 mags brighter than the most luminous WR star in Fig. 8, these are of no use for constraining the progenitor. Furthermore, late time images of NGC 6744 obtained in 2009, when subtracted from the pre-explosion images show no sign of a progenitor disappearance (E. Kankare, private communication). We have hence not considered SN 2005at any further in this work.

4.11 2007gr

SN 2007gr is a Type Ic SN which exploded in NGC 1058; the SN was first discovered by Madison & Li (2007), and spectroscopically classified by Chornock et al. (2007). The photometric evolution of SN 2007gr was similar to that of SN 2002ap (Hunter et al. 2009), but with marked spectroscopic differences. Where spectra of SN 2002ap resembled those of the broad-lined Ic 1998bw, SN 2007gr is spectroscopically similar to a prototypical Type Ic such as SN 1994I. The spectra of SN 2007gr were also notable for the presence of strong carbon absorption (Valenti et al. 2008a). Modelling of late time nebular spectra for SN 2007gr (Mazzali et al. 2010) suggests an ejected oxygen mas of 0.8$M_\odot$, which implies that the CO-core that exploded resulted from a star with ZAMS mass $\sim$15$M_\odot$. Such a progenitor would not have been massive enough to lose its H and He envelopes through stellar winds, and so must have been stripped as part of a binary system.

---

[14] http://www.eso.org/observing/etc/



(a) Pre-explosion WFPC2/WF3 $F450W$ image (b) Pre-explosion WFPC2/PC $F606W$ image (c) Pre-explosion WFPC2/WF3 $F814W$ image (d) Post-explosion ACS/HRC $F814W$ image

Figure 4. Pre-explosion HST+WFPC2 images of the site of SN 2005V, plus a post-explosion HST+ACS image. Each panel covers a $5'' \times 5''$region. Scale and orientation for each panel is identical to that indicated in the post-explosion image.

It was claimed by Paragi et al. (2010) from radio interferometry that SN 2007gr was expanding at relativistic velocities. Together with the absence of extremely high velocities in optical spectra, and the polarization seen in spectropolarimetry (Tanaka et al. 2008), Paragi et al. explained the properties of SN 2007gr with a small amount of material in a low-energy, bipolar jet (seen in radio), coupled with mildly aspherical, non-relativistic ejecta. However, this interpretation was questioned by Soderberg et al. (2010), who claimed from X-ray and radio data that they could fit the evolution of the SN at these wavelengths with a non-relativistic ejecta.

Crockett et al. (2008) used pre-explosion HST and post-explosion Gemini images to show that the SN exploded on the edge of what appears to be either a small compact cluster or a bright supergiant star. Assuming it to be a cluster and fitting of the cluster age yielded uncertain results; Starburst99 models give an age of 7 ± 0.5Myrs or 16 to 21 Myrs with progenitor masses of $28{\pm}4M_{\odot}$ and 14 to $11M_{\odot}$ respectively (Crockett et al. 2008). Regardless of whether the nearby source is a star or a cluster, Crockett et al. (2008b) estimated detection limits for an unseen, point-like progenitor; $F450W > 23.7$ and $F814W > 21.7$. These are not as deep as typical HST WFPC2 images due to the proximity of the SN to this bright object. Assuming $E(B-V) = 0.08$ and a distance of 10.6 Mpc as adopted in Crockett et al. (2008), this corresponds to absolute magnitude limits for the unseen progenitor of $F450W > -6.7$ and $F814W > -8.6$.

4.12 2010br

SN 2010br was discovered in NGC 4051 by amateur astronomers (Nevski et al. 2010) and spectroscopically classified as a Type Ibc SN (Maxwell et al. 2010). NGC 4051 has a recessional velocity distance of 12.7 Mpc (via NED), and a Tully-Fischer distance of 12.2 Mpc (Tully et al. 2009). We adopt the average of these two values, 12.5±0.3 Mpc, as the distance to NGC 4051. The foreground extinction towards SN 2010br is A_V=0.04 mag, from NED, which we adopt as a lower limit to the extinction.

HST pre-explosion imaging consisted of a single WFPC2 F606W image taken on 1994 June 06 (u2e64a01t_c0f). Unfortunately the image was not taken as a cr-split pair, so we have been careful to ensure that there are no cosmic rays in the image at the position of SN 2010br. The SN fell on the PC chip, and so we aligned the PC chip only to a 2340s post-explosion image taken in the F547M filter with WFC3 on 2010 July 17. Using 16 sources we obtained an rms error of 0.2 pixels (10 mas) in a general fit. The SN position was measured in the post-explosion image (ib5e04010_drz) to be 2002.08, 2067.76, with an rms error of $\ll$1 mas. The SN position was transformed to the pre-explosion image, where it was found to be 670.77, 724.56 on the PC chip. We show the images in Figure 5.

No source was detected by hstphot when it was ran on the pre-explosion image with a 3σ threshold. Using the same method as previous, we calculate a limiting magnitude of $F606W > 25.7$ for the progenitor of SN 2010br.

4.13 2011am

Bock, Parker & Brimacombe (2011) discovered SN 2011am in NGC 4219, and Morrell, Stritzinger & Ho (2011) classified the SN shortly after as a young Type Ib SN. Morrell et al. noted the presence of strong NaID absorption at the redshift of the host, suggesting significant extinction. Tully et al. (2009) gives a Tully-Fisher distance for the host of 21.0, while the kinematic distance from NED (corrected for infall on Virgo, GA and Shapely) is 22.5 Mpc. We adopt the Tully-Fisher distance.

Pre-explosion imaging of the site of SN 2011am consists of a 600s WFPC2 exposure in the F606W filter, obtained on 1996 March 16. The SN position lies close to the edge of the PC chip. We attempted to obtain an adaptive optics image of SN 2011am with the VLT+NaCo, however the correction obtained was unsatisfactory due to poor conditions at the time of the observations. Instead, we used an R-band image obtained of SN 2011am on 2011 Mar 27 with the NTT+EFOSC2 under good ($\sim$0.8$''$) seeing conditions to identify the progenitor. The EFOSC2 image was aligned to a pre-explosion Hubble Legacy Archive[15] (HLA) mosaic (u3321601b_drz.fits) to an accuracy of 75 mas. The HLA mosaic was then in turn aligned to the PC chip image, where the transformed SN position was found to be 724.10,355.78, with an uncertainty of 76 mas.

As shown in Figure 6 SN 2011am is close (but probably

---

[15] `http://hla.stsci.edu/`



(a) Pre-explosion WFPC2/PC $F606W$ image



(b) Post-explosion WFC3 $F547M$ image

Figure 5. Pre-explosion HST+WFPC2 image of the site of SN 2010br, plus post-explosion HST+WFC3 image. Each panel covers a $5'' \times 5''$region. Scale and orientation for each panel is identical to that indicated in the post-explosion image, the circle in the upper right corresponds to 5 times the positional rms error.



Figure 6. Pre-explosion HST+WFPC2 image of the site of SN 2011am. The panel covers a $5'' \times 5''$ region, scale and orientation are as indicated. The circle in the upper right corresponds to the positional rms error.

not coincident with) a source with $F606W = 24.204 \pm 0.003$. This source is also classified by HSTPHOT as extended. The formal 5σ limit for the progenitor is 25.8 mag. The extinction towards SN2011am is uncertain but is almost certainly significant. The spectrum at around -6d from Morrell et al. is significantly redder than that of SN2007Y at a similar epoch (-7 days from Stritzinger et al. 2009). We de-reddened the spectrum of SN2007Y by $E(B-V) = 0.112$ as estimated by Stritzinger et al. (2009), and de-reddened the spectrum of SN2011am to match the blue continuum of SN2007Y, finding a very good match with $E(B-V) = 0.6 \pm 0.1$. The intermediate resolution spectrum of Morrell et al. nicely resolves the Na D1 and D2 lines in the Milky Way and NGC4219. The Milky Way components equivalent widths for D1 (5897Å) and D2 (5891Å) are 0.34Å and 0.39Å respectively. Using the Poznanski, Prochaska, & Bloom (2012) relation this suggests $E(B-V) = 0.1 \pm 0.02$, which is in reasonable agreement with the dust map line of sight estimate of Schlegel, Finkbeiner & Davis (1998) $E(B-V) = 0.13$. However there is very strong absorption in the ISM of NGC4219, giving equivalent widths for D1 (5897Å) and D2 (5891Å) of 0.86 and 1.26Å. The Poznanski, Prochaska, & Bloom (2012) relation saturates at equivalent widths of around 0.7Å, hence can't be reliably used and we note that applying it would give values of $E(B-V) = 2.3 - 6.5$, both of which would give unphysical blue spectral slopes and unrealistic absolute magnitudes. Hence we adopt $E(B-V) = 0.6 \pm 0.1$. The corresponding absolute magnitude is given in Table 4

4.14 2011hp

SN 2011hp exploded in the same galaxy as SN 2011am, and so we adopt the same distance of 21.0 Mpc (Tully et al. 2009). The SN was first discovered by Monard (2011a), and classified by Strizinger & Foley (2011) as a Type Ic SN discovered around maximum light. Fraser et al. (2011b) reported a limit for the progenitor of F606W >26.

The pre-explosion image for SN 2011hp is the same as that used for SN 2011am. Unfortunately we have no high resolution imaging for SN 2011hp, and so we are forced to rely on the same natural seeing image as used by Fraser et al. (2011b) for a progenitor identification. We confirm the position measured by Fraser et al. in the mosaic of all four WFPC2 detectors, where the SN falls on the WF4 chip, as shown in Fig. 7. However, we find a shallower limit than that reported by Fraser et al., and calculate a new 5σ limit for the progenitor of SN 2011hp of $F606W > 25.5$.

The spectra of 2011hp are also redder than normal, indicating significant line of sight extinction similar to SN 2011am. We used a spectrum from the ESO programme 184.D-1140 with EFOSC2 on the NTT (3500-9000Å, resolution 17.7Å) taken at +15d after peak (Elias-Rosa et al., in prep) to compare with the low extinction type Ic 2007gr. The spectrum of 2011hp needs to be dereddened by $E(B-V) \simeq 0.5$ to match the spectral shape of 2007gr (when it is dereddened by $E(B-V) = 0.092$). Hence we adopt $E(B-V) = 0.5$, resulting in the absolute magnitude limit reported in Table 4.

4.15 2012au

SN 2012au was identified in the Catalina Sky Survey, and subsequently classified as a Type Ib SN (Silverman et al. 2012). The host galaxy, NGC 4790, has a recessional velocity distance (Virgo corrected) of 21.8 Mpc, and a Tully-Fischer distance of 23.6 Mpc (both from NED). We adopt the latter, corresponding to μ=31.9, and a foreground extinction of $A_V = 0.13$. Pre-explosion imaging consists of 2×160s images in each of $F450W$, $F606W$ and $F814W$, taken with HST+WFPC2. We did not obtain an AO image of SN 2012au, however Van Dyk et al. (2012) set 3σ upper limits of $F450W > 24.8$, $F606W > 25.8$ and $F814W > 25.1$ from an alignment with a natural seeing image. Van Dyk et al. also identified a source slightly outside the transformed SN position (offset by $0.2''$) in the F450W-filter image, with a magnitude of F450W=24.4. This corresponds to an absolute magnitude in B of −7.2 mag. We have re-estimated the limiting magnitudes in these images at the position of the SN and find significantly brighter limiting magnitudes, due to the very high sky background in this vicinity. We determine 5σ limiting magnitudes of F814W> 23.4, F606W> 24.5 and F450W> 24.1.

5 LIMITS FOR THE SAMPLE

Table 4 contains the detection limits for twelve hydrogen deficient core-collapse supernovae. We have not listed metallicity information for these progenitors due to the lack of homogeneous measurements for the host-galaxies. However all the hosts are large spiral galaxies and therefore are expected to have metallicities in the range between that of the Magellanic Clouds and the Milky Way (see discussion in Smartt et al. 2009). Distances in Table 4 are as discussed in the text. To gain insight into the progenitors of Type Ibc SNe, we compare our observed limits to a number of different plausible progenitor populations.

5.1 Limits on progenitors from the observed WR star population

A reasonable hypothesis is that the type Ibc SNe come from massive Wolf Rayet stars. These are stars which will produce Fe-cores and have atmospheres devoid of hydrogen and helium (the WN and WC+WO stars respectively). The classical WR stars are thought to arise in very massive stars that loose their envelopes through stellar winds, potentially enhanced by binary interaction and stellar rotation. These stars appear to be in clusters with turn-off masses that imply lower limits of $> 25 M_\odot$ in the Milky Way and Magellanic Clouds (Massey 2003; Crowther 2007). Due to their high temperatures and high mass-loss rates, the WR stellar radii, masses and bolometric corrections are highly variable. With the strong emission line spectra, this leads to a large range in the absolute magnitudes of the stars. Hence there is no easy way to set meaningful luminosity or mass limits, as can be done for non-detections of II-P progenitor stars (e.g. Smartt et al. 2003; Van Dyk, Li & Filippenko 2003c) and we need a simpler comparison method. If we assume that this population of stars are responsible for the type Ibc SNe that we see locally, we can test if our non-detections are significant. In other words, what is the probability that we have not detected a progenitor star simply by chance? Of course this assumes that the WR stars that we observe now do not change their optical fluxes significantly before core-collapse, and we discuss this debatable point below.



Figure 7. Pre-explosion HST+WFPC2 image of the site of SN 2011hp. The panel covers a $5''\times5''$region, scale and orientation are as indicated. The circle in the upper right corresponds to the positional rms error.

We compare our limits to the observed magnitude distribution of WR stars in the LMC. Massey (2002) presented a catalogue of $UBVR$ photometry for stars in the LMC, and used the WR catalogue of Breysacher, Azzopardi, & Testor (1999) to report broad-band magnitudes for isolated WR stars. This is reasonably complete, and covers a large range of WR luminosities and optical magnitudes. As the catalogue of Massey contains BVR magnitudes for WR stars (whereas most of our progenitor limits are in the HST $F450W$, $F555W$ and $F814W$ filters) we have calculated HST – Johnson-Cousins colours for each of the WR spec-

Table 4. Properties and limits for the observed progenitors in our sample. The broad-band magnitudes with asterisks next to the value are in the standard Johnson-Cousins broad-band filters, the rest are HST filters with: U–$F336W$, B–$F450W$, V–$F555W$, R–$F606W$ and I–$F814W$.

SN	SN Type	Galaxy	Dist(Mpc)	E(B-V)	M_U	M_B	M_V	M_R	M_I	
2000ew	Ic	NGC3810	18.2±3.3	0.04	-	-	-	-6.81	-	
2001B	Ib	IC391	25.5±2.5	0.14	-	-	-8.25	-	-	
2002ap	Ic	NGC628	9.3±1.8	0.09	-8.85*	-4.4*	-	-5.5*	-	
2003jg	Ic	NGC2997	12.5±1.2	0.11	-	-6.30	-	-6.16	-6.99	
2004gn	Ic	NGC4527	14.2±1.3	0.07	-	-	-	-5.55	-	
2004gt	Ibc	NGC4038	22±3	0.07	-9.15	-7.44	-6.07	-	-7.49*	
2005V	Ibc	NGC2146	17.1±3.2	0.90	-	-10.60	-	-8.80	-9.93	
2007gr	Ic	NGC1048	10.6±1.3	0.08		-6.7	-	-	-8.6	-
2010br	Ibc	NGC4051	12.7±2.0	0.04	-	-	-	-4.92	-	
2011am	Ib	NGC4219	24.5±5.6	0.60	-	-	-	-7.44	-	
2011hp	Ic	NGC4219	24.5±5.6	0.50	-	-	-	-7.46	-	
2012au	Ib	NGC4790	23.6±2.0	0.04	-	-8.0	-	-7.5	-8.6	



Figure 8. Limiting magnitudes for the progenitors of Type Ibc SNe (indicated with arrows), compared to the observed magnitudes of LMC WR stars from Massey (2002). WR star magnitudes have been converted to $F450W$, $F555W$ and $F606W$ (indicated by blue, green and red diamonds or arrows) as described in the text, and corrected for foreground extinction ($E(B-V) = 0.080$) and the distance of the LMC ($\mu - 18.50$ mag). All progenitor limits are in the HST filters, with the exception of those for SN 2002ap, which are in the Johnson-Cousins filter system.

tral types, and used these to convert the BVR magnitudes in the Massey (2002) catalogue to the HST filter system.

We first fitted a function of form $Ax + B$ to the effective temperatures of a sample of Galactic WR stars from Hamann, Gräfener & Liermann (2006) of spectral type WNx; where A and B are fitting coefficients. As there were no WN1 stars in the sample of Harmann et al., we have assumed a temperature of 180kK for this subtype. From this fit, we have estimated a temperature for each of the WN spectral types, as listed in Table 5. Temperatures for the WO and WC stars were taken from Sander, Hamann & Todt (2012). We then downloaded model WR spectra from the Potsdam database (Hamann & Gräfener 2004), [16] appropriate to the temperature of each WR subtype. Models were available for a range of radii at each temperature; we have taken the models with the largest and smallest radii as the extreme cases. For each model, we used the IRAF.SYNPHOT package to perform synthetic photometry on the flux-calibrated model spectrum, and from this calculate a colour as listed in Table 5.

Along with spectral type, the colours of the WR stars depend strongly on radius, as the emission lines which dominate WR spectra are stronger in more compact models. The emission line strength will also depend on the wind velocity of the WR star. As these are largely unquantified uncertainties, we will simply adopt average $F450W$-B, $F555W$-V and $F606W$-R colours for each of the WN, WC and WO types, as listed in Table 5.

We plot the observed sample of WR stars in Figure 8 along with the limits from our progenitor sample in each of the $F450W$, $F555W$ and $F606W$ filters.

It is trivial to calculate the probability of not detecting any Type Ibc progenitor thus far, assuming they are randomly drawn from the population of WR stars shown in Fig. 8. For each progenitor with a limit in B,V or R listed in Table 4, we count the number of WR stars with a magnitude brighter than this limit. For most of our Type Ibc SN sample, less than ∼10 per cent of WR stars are brighter than our limit (and hence can be ruled out). We can then calculate the probability of not detecting *any* Type Ibc progenitors to date as $P = \Pi(1 - F_n)$, where F_n is the fraction of WR stars brighter than a particular Type Ibc SN progenitor limit. Using all the limits in Table 4, we find a probability of only 15 per cent that we would not have seen a Type Ibc progenitor thus far.

The WR stars in the LMC which are plotted in Figure 8 are a mix of massive binaries and single stars, while it is of course a single star that explodes. However this does not matter for our test, as if the typical WR population of the LMC were giving rise to Ibc SNe we should see the progenitor stars or their binary systems.

We have performed a similar probability calculation for the sample of WR stars described by Bibby & Crowther (2012, 2010). Here their sample is biased towards brighter

---

[16] http:www.astro.physik.uni-potsdam.dew̃rhPoWR

Table 5. Adopted temperature scale for WR types, together with colours from synthetic photometry of Potsdam models. For each colour, the two values listed are the colour as calculated from the model at that temperature with the smallest and largest radii respectively.

WR Type	Models (R_{min}—R_{max})	$T_{\rm eff}$(kK)	$F450W$-B	$F555W$-V	$F606W$-R	$F814W$-I
WN9	03-10 — 03-04	30	0.026 — 0.042	-0.043 — -0.018	-0.047 — -0.090	-0.102 — -0.090
WN8	04-15 — 04-04	35	0.000 — 0.041	-0.074 — -0.022	0.016 — -0.086	-0.127 — -0.085
WN7	05-16 — 05-04	40	-0.002 — 0.041	-0.081 — -0.030	0.020 — -0.090	-0.123 — -0.081
WN6	07-18 — 07-04	50	-0.006 — 0.047	-0.090 — -0.032	0.024 — -0.092	-0.113 — -0.083
WN5	09-20 — 09-04	60	0.004 — 0.051	-0.074 — -0.029	0.035 — -0.094	-0.119 — -0.044
WN4	11-21 — 11-06	75	0.003 — 0.045	-0.088 — -0.038	0.019 — -0.096	-0.078 — -0.042
WN3	13-21 — 13-08	100	-0.001 — 0.039	-0.133 — -0.040	-0.01 — -0.086	-0.079 — -0.026
WN2	17-21 — 17-12	150	-0.069 — 0.037	-0.207 — -0.042	-0.03 — -0.08	-0.139 — -0.025
WN1	18-21 — 18-13	180	-0.083 — 0.032	-0.226 — -0.043	-0.031 — -0.076	-0.096 — -0.023
WN	mid-range		-0.016	-0.122	-0.030	-0.162
WC5	11-18 — 11-06	80	0.041 — 0.045	-0.059 — -0.034	0.001 — -0.095	-0.879 — -0.875
WC4	15-22 — 15-09	120	0.010 — 0.040	-0.099 — -0.042	0.019 — -0.121	-0.712 — -0.648
WC	mid-range		0.028	-0.071	-0.051	-0.764
WO3	19-26 — 19-24	200	0.004 — -0.064	-0.102 -0.204	0.037 — 0.168	-0.727 — -0.667
WO	mid-range		-0.03	-0.153	0.103	-0.697

magnitudes with no stars in their sample below a $M_V \simeq -4$. If we assume this is representative of the progenitor population then the probabilities in Table 6 decrease. For example, our most restrictive event SN 2002ap would have a probability of 0.01. The probability for the entire population would be 4×10^{-6} and so essentially zero, even without this SN. This demonstrates how difficult it is to detect Wolf-Rayet stars in other galaxies at great distances let alone progenitors that may also be fainter than these Wolf-Rayet stars. It also demonstrates that this bright end of the WR luminosity function is almost certainly not producing the Ibc SNe that we see in the local Universe.

5.2 Limits on progenitors from population modelling

Comparing the progenitor limits from archival imaging to the observed population of Wolf-Rayet stars is the simplest approach to constraining the progenitor population of Type Ibc SNe. However, it is limited by the fact that observed Wolf-Rayet stars will lie within a range of different evolutionary states, with some stars closer to core-collapse than others. It is possible that Type Ibc progenitors become fainter in optical bands as they evolve closer to the point of core-collapse (e.g. Yoon et al. 2012). To account for this we compare our observed progenitor limits to synthetic progenitor populations created with BPASS. The four synthetic populations we use are as follows:

(i) Single WR stars with $X_{\rm surface} < 0.001$ and $\log(T_{\rm eff}/{\rm K}) > 4$

(ii) Binary WR stars with $X_{\rm surface} < 0.001$, $\log(T_{\rm eff}/{\rm K}) > 4$ and $M > 2M_{\odot}$

(iii) Single star Type Ibc progenitors with $X_{\rm surface} < 0.001$, $\log(T_{\rm eff}/{\rm K}) > 4$ and have completed core helium burning

(iv) Binary type Ibc progenitors with $X_{\rm surface} < 0.001$, $\log(T_{\rm eff}/{\rm K}) > 4$, $M > 2M_{\odot}$ and have completed core helium burning

The SN types resulting from the stars in these synthetic populations are determined using the parameters listed in table 2. For the binary population we include the contribution of the primary and secondary stars to the WR population. We also consider the companion flux when calculating the observed magnitude of the system. We see in Figure 9 that the synthetic single-star WR population has a similar mean magnitude to the observed LMC WR population used above, but is not able to reproduce the spread of magnitudes. In contrast, the synthetic binary WR population is a much closer match to the observed distribution. This is because binary interactions increase the amount of mass-loss a star can undergo, and so widens the range of initial masses (and hence luminosities) which give rise to WR stars. In addition the inclusion of the flux from the secondary in binary systems leads to some WR stars being apparently more luminous than possible from single stars alone.

We stress that our method differs from that of Yoon et al. (2012), who consider the very end point of each evolutionary track as the point of explosion. We assume that our model stars can explode at any time after the end of core-helium burning. The advantage of this is that it gives us a range of possible magnitudes for each model, and hence allows us to include the uncertainties of stellar evolution and Wolf-Rayet atmosphere modelling in our analysis. The disadvantage is that we may overestimate the magnitude of single-star progenitor models, which we find to be higher than those suggested by Yoon et al. (2012). Furthermore they estimate pre-SN magnitudes in the narrow band $M_{\rm v}$ rather than the broad-band filter (M_V) commonly available for progenitor detections. The narrow band filter does not include emission line flux which can contribute significantly to the luminosity of Wolf-Rayet stars, and $M_{\rm v}$ can be typically 0.5-1.0 magnitudes fainter than broadband BVR. These factors would lead in turn to an increase in the probability that no progenitor would have been observed in pre-explosion images. This can be seen in Table 6, as the probability of a detection for our single-star WR population is lower than that for our single-star progenitor population. However the difference in luminosity may be more extreme then we provide here, and this caveat must be considered when comparing the probabilities of the different synthetic populations.

Using these synthetic populations we can again compare the progenitor limits to the various synthetic populations. From each SN we can calculate a probability that the progenitor would remain undetected from the fraction of the synthetic population with a magnitude less than the limit. We then combine those individual probabilities to estimate the probability that no progenitor would have been observed in any of the pre-explosion images. We show the resulting probabilities and their combination in Table 6. As mentioned previously, the probability that we would not have detected a Type Ibc progenitor if the LMC WR sample is representative of the progenitors is 0.15. Similarly low values are found for all the progenitor models, although the population of synthetic binary progenitors has a slightly higher probability at 0.27.

We note that without the deep limit for SN 2002ap, in all cases the probability that we would not have observed a SN progenitor increases significantly. However, the probability of the binary population do not increase by as much as that of the single star models. This is because the range of possible magnitudes in a binary is greater than those from single stars. While this means there are more low luminosity progenitors, there are also more progenitors at the high luminosity end of the distribution. This gives the false impression that single stars are more likely to be the possible progenitors.

In conclusion, while we favour a mixed population of Wolf-Rayet stars and lower mass helium stars in binaries as the progenitors of Type Ibc SNe, we cannot set strong constraints from the current limits and non-detections. Furthermore, our limits on the progenitors are constrained by the depth of the pre-explosion images, and strongly dependent on the deep limits for two of our sample, namely SNe 2002ap and 2010br.

6 DISCUSSION

There are two main candidates for the progenitors of Type Ibc SNe: Wolf-Rayet stars ($M \gtrsim 20$ $M_\odot$) and lower mass helium-stars. The former will experience mass loss which is primarily due to stellar winds, and will effectively evolve as single stars. The latter are the result of close binary evolution, where the mass-loss is due to binary interactions of Roche-lobe overflow or common envelope evolution. Both progenitor channels are plausible. WR stars are commonly observed in star-forming environments, but to the best of our knowledge no completely hydrogen-free low-mass helium stars are known (with a core mass that will create an ONeMg or Fe-core and experience core-collapse) and their existence remains conjecture.



Figure 9. Cumulative frequency $F555W$ magnitude distributions for observed WR stars (black line), synthetic single WR stars (blue-dashed line), synthetic single star Type Ibc progenitors (blue-solid line), synthetic binary WR stars (red-dashed line), synthetic binary Type Ibc progenitors (red-solid line)

6.1 Arguments in favour of interacting binaries as progenitors of Ibc SNe

From the limits presented in Section 5, if Wolf-Rayet stars were the sole progenitors of Type Ibc SNe it is likely that we would have observed one as a progenitor in pre-explosion imaging. The number of non-detections to date suggest that at least some, and possibly most Type Ibc SNe arise from lower mass stars in binary systems. This is further supported by the fact the binary models can reproduce the observed relative rate of different SN types (Podsiadlowski et al. 1992; Eldridge, Izzard & Tout 2008; Yoon, Woosley & Langer 2010; Eldridge, Langer & Tout 2011). Current single star models predict too many Type II SNe, however recent studies have indicated that the mass-loss rates of red supergiants remain a large source of uncertainty (Yoon & Cantiello 2010; Georgy 2012).

Currently no single star model can fit the population of RSGs, WR stars and SNe rates simultaneously. The Geneva rotating stellar evolution models have come close (Ekström et al 2012), but assume a single initial rotational velocity rather than a distribution of rotation velocities as are observed (Hunter et al. 2008). In contrast, binary models can also come close to reproducing both observed stellar populations and relative SN rates with reasonable distribution of initial binary parameters (Eldridge, Izzard & Tout 2008; Eldridge, Langer & Tout 2011). In our binary population model which fits best both the observed relative SN rates and the non-detection of a Type Ibc progenitor, we find that single WR stars contribute one-fifth of Ibc SNe, with binary systems giving rise to the rest. Furthermore in our synthetic population only half of the SNe will be in a binary at the time of core-collapse, the remainder being apparently single. This is because binaries are either unbound in the first SN or have a compact companion at the time of the second SN. Deep post-explosion images with the *Hubble Space Telescope* at the sites of Type Ibc SNe may help identify the surviving binary companion of the progenitor.

It is important to note that our binary population model also predicts the location on the HR diagram of the progenitors of hydrogen-poor Type IIb and IIL SNe. In Fig-

Table 6. The probabilities that no progenitor would have been observed given the different models.

			Progenitor models			
SN	SN Type	Observed WRs	Single WR stars	Single Progenitors	Binary WR stars	Binary progenitors
2000ew	Ic	0.96	1.00	1.00	0.94	0.97
2001B	Ib	0.99	1.00	1.00	1.00	1.00
2002ap	Ic	0.48	0.08	0.16	0.21	0.49
2003jg	Ibc	0.93	1.00	1.00	0.87	0.94
2004gn	Ic	0.75	0.94	0.99	0.76	0.90
2004gt	Ibc	0.94	0.98	0.99	0.83	0.91
2005V	Ibc	0.99	1.00	1.00	1.00	1.00
2007gr	Ic	0.95	1.00	1.00	0.90	0.95
2010br	Ibc	0.57	0.62	0.76	0.56	0.81
2011am	Ib	0.98	1.00	1.00	0.98	0.99
2011hp	Ib	0.98	1.00	1.00	0.98	0.99
2012au	Ib	1.00	1.00	1.00	0.98	0.99
Model Probability		0.15	0.05	0.12	0.05	0.27

ure 11 we show a cartoon HR diagram based on our synthetic population, where the YSG progenitors lie above our predicted location for the hydrogen-deficient SNe. We note that this matches the observed progenitors of SNe 1993J, 2008ax, 2009kr and 2011dh, which are intermediate between the Type Ibc progenitors and hydrogen rich IIP RSG progenitors.

Hence the question then is, where are the binary systems that harbour Ibc progenitor systems? If we examine our binary evolution models more closely we find that stars below approximately $15M_\odot$ spend only a very small fraction of their total lifetime without hydrogen in their structure before core collapse. This is shown in Figure 10 which displays the hydrogen mass fraction of single-star and binary models before the end of the stellar models. Stars less massive than $15M_\odot$ retain a low mass hydrogen envelope that is helium rich. This is only lost a short time before core collapse. In the right-hand panel of Figure 11 the envelope is lost as the stars evolve from their hottest extent to cooler effective temperatures. Stars with parameters similar to these stars have been observed and are objects such as V Sagittae, WR7a and HD45166[17] (Steiner & Diaz 1998; Oliveira et al. 2003; Steiner & Oliveira 2005; Groh, Oliveira & Steiner 2008). The objects are often discarded as potential Ibc progenitors due to the presence of hydrogen on their surface. However in Figure 10 we show the surface hydrogen mass fraction for our models which indicate that these possible type Ibc progenitors are hiding large helium cores and the residual hydrogen could be stripped in the last 10^4 yrs of their life. Hence it may not be surprising that they exist with in this configuration and that completely hydrogen free He-stars in binaries are difficult to locate.



Figure 10. Diagram of the hydrogen mass fraction of different progenitor models before the end of our models. The solid and dotted blue lines are for $30M_\odot$ and $50M_\odot$ single-star models respectively. The solid and dotted black lines are for $15M_\odot$ and $20M_\odot$ binary models respectively. The solid and dotted red lines are for $10M_\odot$ and $12M_\odot$ binary models respectively. .

This may seem unexpected but we stress that a binary interaction does not immediately lead to complete loss of the hydrogen envelope in a star. Roche-Lobe overflow or common envelope evolution removes most of the hydrogen envelope but eventually an envelope mass is reached at which point it collapses back within the star's Roche Lobe. Stellar winds are then responsible for removing the remaining gas. For stars above $15M_\odot$ the winds are strong enough to do this during helium burning. Below this limit it is not until the star evolves to a helium giant that the mass-loss rate becomes strong enough to drive off the hydrogen before core-collapse. We note that at lower metallicity weaker winds may reduce the number of type Ibc SNe from these binary stars, therefore the type Ibc rate will decrease at lower metallicities.

The other piece of observational evidence that supports helium stars as SN progenitors is the ejecta masses deter-

---

[17] HD45166 is slightly more hydrogen rich than predicted by our models. This is probably a result of the secondary star filling its Roche lobe at periastron and transfering hydrogen back onto the primary (Steiner & Oliveira 2005).



Figure 11. Cartoon HR diagrams of SN progenitors, the red, yellow and blue regions show the expected location of progenitors for type IIP, other type II and type Ib/c SNe. The left panel shows the single-star scenario. The solid lines show the evolution tracks for stars with masses given at their initial location. The right panel shows the binary scenario with the solid tracks at 10, 15 and 20M$_\odot$ showing binary evolution tracks and the dashed lines the single star tracks. In both plots the points with error bars show the locations of SNe 1987A, 1993J, 2008cn, 2009kr and 2011dh (Podsiadlowski 1992; Maund et al. 2004; Elias-Rosa et al. 2009a, 2010; Fraser et al. 2010b; Maund et al. 2011) respectively. The circles show the progenitor locations of observed type IIP progenitors and the arrows the upper limits for these progenitors (Smartt et al. 2009).

mined from the studies of the lightcurves and observed ejecta velocities. There are now many Ibc SNe with well sampled lightcurves, and some with data stretching through the nebular phase. Simple analytical models following Arnett (1982) can estimate ejecta mass and ^{56}Ni mass from the rise time, width and peak of the lightcurve. More sophisticated radiative transfer modelling with Monte Carlo techniques (for example Nakamura et al. 2001; Mazzali et al. 2007; Dessart et al. 2011) have been used to determine also the kinetic energy from the velocity information in the spectra. A summary of the main results for type Ibc SNe is compiled in Table 7. The models of the most common Ibc SNe types in the local Universe (within our 28 Mpc volume), and those for which we have limited the progenitor, all tend to be between 1-3M$_\odot$. This fact has recently been highlighted by sophisticated modelling of Ibc SNe by Hachinger et al. (2012) and Dessart et al. (2012). In addition, Drout et al. (2011) presented a homogeneous study of the ejecta masses of twenty-five Ibc SNe and found similarly low masses consistently through the sample. There are a small percentage of the overall Ic SNe class that have very broad lightcurves and usually, but not always, high ejecta velocities (often called Ic-BL, or broad-lined Ic). There is some evidence that they have significantly higher ejecta masses. Their lightcurves are significantly broader and a higher ejecta mass is the obvious way to explain the differences with a similar physical model. However these Ic-BL are rare, with only one (SN 2002ap) having been discovered within 28 Mpc in the duration of our survey. Their contribution to the overall Ibc SN population is likely less than 3 per cent (see Section 2) and these could conceivably be from higher mass He or CO stars. Wolf-Rayet stars would be the obvious candidates for some of these type Ic SNe (Dessart et al. 2012).

In Table 7 we also list the oxygen ejecta mass where it has been estimated by model application. Generally this is around 1M$_\odot$ or lower, also supporting the idea that the bulk of the Ibc SNe are not from WR stars progenitors with zero-age main-sequence masses $\geqslant 25$M$_\odot$ (see Mazzali et al. 2010).

We predict ejecta masses from the synthetic populations as described in Eldridge & Tout (2004b). The should be considered upper limits as we do not evolve our models to Fe-core formation and so may underestimate the final remnant mass. Our binary progenitor models (which mainly form neutron-star remnants) have an average ejecta mass of 4.2 ± 2.4M$_\odot$ while single star progenitors (which mainly form black-hole remnants), have ejecta masses in the range to 6.6 ± 0.9M$_\odot$. This further supports the idea that *normal* type Ibc SNe come from helium stars formed in binaries with initial masses below 20M$_\odot$ and form neutron stars at core collapse.

6.2 Arguments in favour of Wolf-Rayet stars as progenitors of Ibc SNe

Recently, much work has been done on the stellar populations near the location of SNe (e.g. Murphy et al. 2011; Leloudas et al. 2011; Anderson et al. 2012). These range from studying the resolved stellar populations using high resolution HST data, such as that presented here

Table 7. Measured ejecta masses for Ibc SNe from the literature. These are lightcurve modelling. In some cases an estimated mass of oxygen is also determined from spectral modelling and is added here for reference. Typical stellar evolution models predict that a 25$M_\odot$ star would produce between 2-4$M_\odot$ of oxygen during its nucleosynthetic evolution.

SN	Type	O mass / $M_\odot$	Total ejecta mass / $M_\odot$	Refs
2007Y	Ib	0.2	0.4	Stritzinger et al. (2009)
1994I	Ic	..	0.7	Valenti et al. (2008b)
2007gr	Ic	0.8	1-2	Mazzali et al. (2010), Hunter et al. (2009)
2006aj	Ic	1.3	2	Mazzali et al. (2006), Maeda et al. (2007)
2002ap	Ic	1.2	2.5	Mazzali et al. (2007)
2003jd	Ic	...	3	Valenti et al. (2008b)
2008D	Ib	1.1	3.6	Dessart et al. (2012)
2004aw	Ic	...	5	Valenti et al. (2008b)
2009jf	Ib	...	5-7	Valenti et al. (2011c)
1997ef	Ic	...	7-10	Mazzali, Iwamoto, & Nomoto (2000)
1998bw	Ic	2-3	10	Nakamura et al. (2001)
2011bm	Ic	5-10	7-17	Valenti et al. (2012)
Sample Average	Ib	...	$2^{+1.1}_{-0.8}$	Drout et al. (2011)
Sample Average	Ic	...	$1.7^{+1.4}_{-0.9}$	Drout et al. (2011)
Sample Average	Ic-BL	...	$4.7^{+2.3}_{-1.8}$	Drout et al. (2011)

(Murphy et al. 2011) to lower resolution ground-based data that typically use Hα to probe regions of star formation. Using the latter technique Anderson et al. (2012) propose that type Ibc SNe tend to show stronger spatial association with Hα emission in galaxies compared to type II SNe, and type II-P in particular. As Hα emission originates in young clusters and associations, their interpretation is that type Ibc SNe may arise from significantly higher mass stars than type II-P. They also find that the spatial correlation is strongest for type Ic SNe, even more so than type Ib. Anderson et al. (2012) interpret this as an increasing mass sequence, with type II SNe coming from the lowest mass stars and Ib and Ic from progressively higher mass. This would argue in favour of Ibc progenitors being significantly higher mass stars than progenitors of II-P SNe, and since we know that II-P SNe arise in 8-17$M_\odot$ stars (Smartt et al. 2009; Smartt 2009) then this might suggest WR progenitors of type Ibc SNe with initial mass $\geqslant$20$M_\odot$.

These SNe are typically at distances of 50-100 Mpc and the ground-based imaging does not specifically identify the host HII region, a fact which led Crowther (2012) to study the environments of the closest 39 CCSNe in nearby ($\leqslant$15 Mpc) and low inclination galaxies. While Ibc SNe are nearly twice as likely to be associated with nearby H II emission, Crowther (2012) finds that these nebulae are long lived, giant H II regions with typical lifetimes more than 20 Myrs. Hence the association only provides weak constraints on Ibc progenitor masses. Binary evolution would extend the period during which Type Ibc SNe occur up to 20 Myrs and this is still roughly in agreement with observed population ages from the H II region analysis. The greater mean age of binary SN progenitors is also in accord with the absence of any HII regions or large young stellar populations at the site of the Type Ibc SNe in our sample. The typical lifetime of a HII region is between 3 to 5 Myrs, i.e. shorter than the lifetime of most binary Type Ibc progenitors. Only the most massive regions such as NGC604 or 30 Doradus have more extended periods of star-formation, and such systems are rare.

There are two other factors that may have prevented us from detecting WR progenitor stars. Firstly it is possible that WR stars evolve rapidly in the latter stages of their lives. This could mean that, just prior to core-collapse, they appear significantly different to the WR stars we observe in the LMC, possibly getting hotter and fainter in the last 5 per cent of their lifetimes. This has been investigated by Yoon et al. (2012). However we suggest that progenitors are slightly more observable than in the work of Yoon et al. (2012). They determined magnitudes in the narrow band $M_{\rm v}$ rather than the broad-band filter commonly available for progenitor detections. The narrow band filter does not include emission line flux which can contribute significantly to the luminosity of Wolf-Rayet stars, and $M_{\rm v}$ can be typically 0.5-1.0 magnitudes fainter than broadband BVR.

Secondly, we may have underestimated the extinctions towards the WR star progenitors (and hence our detection limits are not as deep as we have presented here), or WR stars could create and eject dust in the last fraction of their lifetimes. The values for $E(B-V)$ used in Table 4 are those estimated for the foreground line of sight (including host galaxy extinction where possible). The distribution is characterised by a mean and standard deviation of $< E(B-V) >= 0.22 \pm 0.28$, which is similar to the extinctions estimated by Bibby & Crowther (2012, 2010) towards WR stars in NGC5068 and NGC7793. In addition, we have added our estimates of extinction to the limiting magnitudes, whereas for the LMC sample we have simply taken foreground Milky Way extinction. Any extra internal LMC, or circumstellar extinction would tend to make the LMC reference sample brighter for our comparison purposes and the restrictions tighter. In this study we have no ability to disentangle the foreground extinction from circumstellar, hence we have not employed the new and physically consistent methods of Kochanek, Khan, & Dai (2012). If a Ibc progenitor were found then it would be necessary to consider the circumstellar extinction separately to foreground ISM as discussed by Kochanek, Khan, & Dai (2012) as is the case for SN2012aw.

7 CONCLUSIONS

We have used the observed limits on the progenitors of Type Ibc SNe in pre-explosion images, the observed relative rates for different SN types and a binary population model to study their progenitor population. From both sets of data, we find that a single-stellar population is disfavoured, and that the bulk of the type Ibc SNe most likely arise from low mass progenitors which have been stripped by a binary companion. This scenario is also supported by the fact that the ejecta masses from Ibc SNe are, in general, too low to be from massive Wolf-Rayet stars.

The favoured progenitors are therefore the hypothetical low-mass helium stars. However no such hydrogen-free stars with masses large enough to experience core-collapse are known in the Galaxy. Our stellar models of these systems indicate that this is actually to be expected. The binary interaction that seals these stars to their final fate does not remove all hydrogen from the stellar surface and leaves approximately a few times $0.1\mathrm{M}_{\odot}$. This is removed by stellar winds as the star evolves at the end of core helium burning. These objects would appear similar to observed binary systems such as V Sagittae, WR7a and HD45166. Therefore we suggest that the final fate of these systems will be type Ibc SNe. Further detailed study and modelling of these systems are required to confirm this.

While there are some Ic SNe which have broad lightcurves and likely large ejecta masses, they are much rarer by volume compared to the bulk of the normal Ibc SN population. It's still possible that these Ic SNe do come from massive Wolf-Rayet stars, however Podsiadlowski et al. (2004) show that even if one assumes that only stars above $80\mathrm{M}_{\odot}$ produce broad-lined and broad lightcurve Ic SNe then the observed rate of such SNe is still too low. Hence the massive, classical Wolf-Rayet population does not produce the majority of the normal Ibc SNe we see within 28 Mpc. However they could contribute to this population or supply progenitor systems producing rarer type Ibc events. It's possible that some fraction of WR stars form black holes in such a way that they produce faint SNe or no visible display.

8 ACKNOWLEDGEMENTS

The research leading to these results has received funding from the European Research Council under the European Union's Seventh Framework Programme (FP7/2007-2013)/ERC Grant agreement n° [291222] (PI : S. J. Smartt). We thank Erkki Kankare for information on SN 2005at. We thank Nancy Elias-Rosa and Max Strizinger for sharing spectra of 2011br and 2011am. We thank Brad Cenko for information on SN 2004gn. We thank Stefano Benetti and Nancy Elias Rosa for access to an NTT image of SN2011am and a spectrum of SN2011hp both from the ESO Large Programme 184.D-1140 'Supernova Variety and Nucleosynthesis Yields'

REFERENCES

Agnoletto I., Harutyunyan A., Benetti S., Turatto M., Cappellaro E., Dennefeld M., Adami C., 2007, CBET, 1163, 1
Aldering G., Humphreys R.M., Richmond M., 1994, AJ, 107, 662
Aldering G. et al., 2006, ATEL, 690
Anderson J., Morrell N., Folatelli G., Stritzinger M., 2009, CBET,1789, 1
Anderson J.P., Habergham S.M., James P.A., Hamuy M., 2012, MNRAS, 424, 1372A
Arcavi I. et al., 2011, ApJ, 742, 18
Arnett W. D., 1982, ApJ, 253, 785
Ayani K., Yamaoka H., 1999, IAUC, 7137, 1
Ayani K., Kawabata T., Ioroi M., Ohshima O., 2001, IAUC, 7683
Ayani K., Kawabata T., Yamaoka H., 2002, IAUC, 7864, 4
Ayani K., Yamoka H., 2002, IAUC, 7976, 3
Ayani K., Hashimoto T., Yamaoka H., 2003, IAUC, 8048, 2
Balam D.D., Graham M.L., Hsiao E.Y., 2011, CBET, 2625, 3
Begelman M.C., Sarazin C.L., 1986, ApJ, 302, L59
Benetti S., et al. 2004, MNRAS, 348, 261
Benetti S., Valenti S., Magazzu A., Harutyunyan A., 2009, CBET 1667, 1
Berger E., 2010, ATEL, 2655
Bessell M.S., Schmidt B.P., 2010, 2337, 1
Bibby J. L., Crowther P. A., 2010, MNRAS, 405, 2737
Bibby J. L., Crowther P. A., 2012, MNRAS, 420, 3091
Binder B., Williams B.F., Kong A.K.H., et al. 2011, ApJ, 739, L51
Blackman J., Schmidt B., Kerzendorf W., 2006, CBET, 541, 1
Blondin S., Masters K., Modjaz M., Kirshner R., Challis P., Matheson T., Berlind P., 2006, CBET, 636, 1
Blondin S., Calkins M., 2007a, CBET, 800, 2
Blondin S., Calkins M., 2007b, CBET, 1119, 1
Bock G., Parker S., Brimacombe J., 2011, CBET, 2667, 1
Bond H.E. et al. 2003, Nature, 422, 405
Borsato, L., Nascimbeni, V., Benetti, S., et al. 2012, Central Bureau Electronic Telegrams, 2993, 2
Botticella M.T., Pastorello A., Smartt S.J., et al. 2009, MNRAS, 398, 1041
Botticella M. T., Smartt S. J., Kennicutt R. C., Cappellaro E., Sereno M., Lee J. C., 2012, A&A, 537, A132
Branch D., et al., 2002, ApJ, 566, 100
Branch D., 2003, AJ, 126, 1489
Breysacher J., Azzopardi M., Testor G., 1999, A&AS, 137, 117
Brunthaler A., Menten K.M., Reid M.J., Henkel C., Bower G.C., Falcke H., 2009, A&A, 499, L17
Brunthaler A. et al. 2010, A&A, 516, 27
Burrows A., 2012, arXiv, arXiv:1210.4921
Bufano F., Benetti S., Pastorello A., 2011, CBET, 2676, 1
Challis P., 2008, CBET 1628, 2
Challis P., 2009, CBET 1673
Chen J., Wang., X.-F., Wang X.-L., et al. 2011, Central Bureau Electronic Telegrams, 2943, 2
Chevalier R.A., Soderberg A.M., 2010, ApJ, 711, L40
Chornock R., Filippenko A.V., 2001, IAUC, 7577, 2
Chornock R., Filippenko A.V., Li W., Foley R.J., Reitzel D., Rich R.M., 2007, CBET, 1036, 1
Chornock R., Berger E., 2010, CBET, 2086, 1
Connolly A., 2004, IAUC, 8359, 1

Contreras C., Morrell N., Gonzalez S., Lee K-G., 2007, CBET, 1068, 1

Crockett R.M. et al., 2007, MNRAS, 381, 835

Crockett R., et al., 2008b, ApJ, 672L, 99C

Crockett R.M., Smartt S.J., Pastorello A., Eldridge J.J., Stephens A.W., Maund J.R., Mattila S., 2011, MNRAS, 410, 2767

Crowther P.A., 2007, ARA&A, 45, 177C

Crowther P. A., 2012, MNRAS, 183

Dennefeld M., Patris J., 2000, IAU Circ., 7532, 3

Dessart L. et al., 2008, ApJ, 675, 644

Dessart L., Hillier D.J., Livne E., Yoon S.-C., Woosley S., Waldman R., Langer N., 2011, MNRAS, 414, 2985D

Dessart L., Hillier D. J., Li C., Woosley S., 2012, MNRAS, 424, 2139

Di Carlo E., et al. 2002, ApJ, 573, 144

Drout M.R. et al., 2011, ApJ, 741, 97D

Ekström S. et al., 2012, A&A, 537A, 146E

Eldridge J.J., Tout C.A., 2004b, MNRAS, 353, 87

Eldridge J.J., Izzard R.G., Tout C.A., 2008, MNRAS, 384, 1109E

Eldridge J.J., Stanway E.R., 2009, MNRAS, 400, 1019

Eldridge J.J., Langer N., Tout C.A., 2011, MNRAS, 411, 3501

Elias-Rosa N., et al. 2006, MNRAS, 369, 1880

Elias-Rosa N., et al. 2004a, IAUC, 8273, 2

Elias-Rosa N., et al. 2004b, IAUC, 8376, 2

Elias-Rosa N. et al 2008, MNRAS, 384, 107

Elias-Rosa N., 2009a, ApJ, 706, 1174

Elias-Rosa N., van Dyk S.D., Agnoletto I., Benetti S., 2009c, CBET, 1765, 1

Elias-Rosa N. et al., 2010, ApJL, 714, L254

Elias-Rosa N. et al., 2011, ApJ, 742, 6

Elmhamdi A., Tsvetkov D., Danziger I.J., Kordi A., 2011, ApJ, 731, 129

Fassia A., et al., 2001, MNRAS, 325, 907

Filippenko A.V., 1997, ARA&A, 35, 309F

Filippenko A.V., De Breuck C., 1998a, IAUC, 6994, 3

Filippenko A.V., De Breuck C., 1998b, IAUC, 6997, 2

Filippenko A.V., Leonard D.C., Riess A.G., 1998, IAUC, 7089, 3

Filippenko A.V., Li W.D., Modjaz M., 1999, IAUC, 7152, 2

Filippenko A.V., Foley R.J., Modjaz M., 2000, IAUC, 7547, 2

Filippenko A.V., Chornock R., 2000, IAUC, 7511, 2

Filippenko A.V., Chornock R., 2001, IAUC, 7638, 1

Filippenko A.V., Chornock R., 2003, IAU Circ., 8211, 2

Filippenko A.V., Ganeshalingam M., Serduke F.J.D., Hoffman J.L., 2004, IAUC, 8404, 1

Filippenko A.V., Foley R.J., Chornock R., Matheson T., 2004, IAUC, 8420, 2

Filippenko A.V., Foley R.J., 2005, IAUC, 8486, 3

Filippenko A.V., Silverman J.M., Foley R.J., Modjaz M., Papovich C., Willmer C.N.A., Blondin S., Brown W., 2007, CBET, 1101, 1

Folatelli G., Morrell N., Phillips M., Hamuy M., 2007, 862, 1

Folatelli G., Gonzalez S., Morrell N., 2007, 850, 1

Folatelli G., Olivares F., Morrell N., 2008, 1227, 1

Folatelli G., Morrell N., 2009, CBET, 1807, 1

Foley R.J., Pugh H., Filippenko A.V., 2004, IAUC, 8374, 2

Foley R.J., Wong D.S., Ganeshalingam M., Filippenko A.V., Chornock R., 2004, IAUC, 8339, 2

Foley R.J., Smith N., Ganeshalingam M., Li W., Chornock R., Filippenko A.V., 2007, ApJ, 657, 105

Foley R.J. et al. 2009, AJ, 138, 376

Foley R.J., 2009, CBET, 1858, 1

Foley R.J., Brown P.J., Rest A., Challis P.J., Kirshner R.P., Wood-Vasey W.M., 2010a, ApJ, 708, 61

Foley R.J. et al. 2010b, AJ, 140, 1321

Foley R.J., Berger E. Fox O., Levesque E.M., Challis P.J., Ivans I.I., Rhoads J.E., Soderberg A.M., 2011, ApJ, 732, 32

Foley R.J., Kromer M., Marion G.H., et al. 2012, ApJ, 753, L5

Fraser M., Smartt, S.J., Crockett M., Mattila S., Stephens A., Gal-Yam A., Roth K., 2010a, ATEL, 2131, 1

Fraser M., Smartt S. J., Crockett M., Mattila S., Stephens A. G.-Y. A., Roth K., 2009, ATel, 2131, 1

Fraser M. et al., 2010b, ApJL, 714, L280

Fraser M. et al., 2011, MNRAS, 417, 1417

Fraser M., Maund J.R., Smartt S.J., Bufano F., Benetti S., Elias-Rosa N., 2011b, ATEL, 3371, 1

Fraser, M., Maund, J. R., Smartt, S. J., et al. 2012, ApJ, 759, L13

Fraser M. et al., 2012, in prep

Gal-Yam A., Ofek E.O., Shemmer O., 2002, MNRAS, 332, 73

Gal-Yam A., et al. 2005, ApJ, 630, 29

Gal-Yam A., Cenko S.B., Fox D.B., Leonard D.C., Moon D.-S., Sand D.J., Soderberg A.M., 2007, AIPC, 924, 297

Gal-Yam A., et al. 2008, ApJ, 685, 117

Garnavich P., Jha S., Kirshner R., Challis P., Berlind P., Holman M., IAUC, 7306, 2

Garnavich P.M., et al, 2004. ApJ, 613, 1120

Gaskell C.M., Cappellaro E., Dinerstein H.L., Garnett D.R., Harkness R.P., Wheeler J.C.,1986, ApJ, 306, L77

Georgy C., 2012, A&A, 538L, 8G

Gerardy C.L., Fesen R.A., Nomoto K., Maeda K., Hoflich P., Wheeler J.C., 2002, PASJ, 54, 905

Gerardy C.L. et al. 2007, ApJ, 661, 995

Gómez G., López R., 2000, AJ, 120, 367

Gonzalez L., Morrell N., Hamuy M., 2003, IAUC, 8173, 3

Gonzalez S., Krzeminski W., Folatelli G., Hamuy M., Morrell N., 2004, IAUC, 8409, 2

Graham J., Li W., 2003, IAU Circ., 8207, 1

Groh J.H., Oliveira A.S., Steiner J.E., 2008, A&A, 485, 245

Hachinger S., Mazzali P. A., Taubenberger S., Hillebrandt W., Nomoto K., Sauer D. N., 2012, MNRAS, 422, 70

Harmann W.-R., Gräfener G., 2004, A&A, 427, 697

Harmann W.-R., Graäfener G., Liermann A., 2006, A&A, 457, 1015

Harutyunyan A., Agnoletto I., Benetti S., Turatto M., Cappellaro E., Lorenzi V., 2007a, CBET, 903, 1

Harutyunyan A., Benetti S., Cappellaro E. Patat F., Ghinassi F., 2007b, CBET, 1021, 1

Harutyunyan A., Navasardyan H., Benetti S., Turatto M., Pastorello A., Taubenberger S., 2007c, CBET, 1184, 1

Harutyunyan A.H. et al. 2008, A&A, 488, 383

Harutyunyan A., Bufano F., Benetti S., 2009, CBET, 1722, 1

Hamuy M., Roth M., Morrell N., 2002, IAUC, 8037, 2

Heger A., Fryer C.L., Woosley S.E., Langer N., Hartmann D.H., 2003, ApJ, 591, 288

Hendry M.A., et al. 2005, MNRAS, 359, 906

Hendry M.A. et al., 2006, MNRAS, 369, 1303

Hess K.M., Pisano D.J., Wilcots E.M., Chengalur J.N., 2009, ApJ, 699, 76

Hernandez M. et al., 2000, MNRAS, 319, 223

Hillebrandt W., Niemeyer J.C., 2000, ARA&A, 38, 191

Hoffman D., et al. 2011, ATEL, 3160, 1

Horiuchi S., Beacom J.F., Kochanek C.S., Prieto J.L., Stanek K.Z., Thompson T.A., 2011, ApJ, 738, 154

Howell D.A., 2003, IAUC, 8241, 2

Howerton S. et al. 2011, CBET, 2962, 1

Humphreys R.M., Prieto J.L., Rosenfield P., et al. 2010, ApJ, 718, L43

Hunter I., Lennon D.J., Dufton P.L., Trundle C., Simn-Daz S., Smartt S.J., Ryans R.S.I., Evans C.J., 2008, A&A, 479, 541

Hunter D.J. et al., 2009, A&A, 508, 371

Immler S., et al. 2006, ApJ, 648, L119

Iwamoto K., Nomoto K., Hoflich P., Yamaoka H., Kumagai S., Shigeyama T., 1994, ApJL, 437, L115

Nomoto K. I., Iwamoto K.,Suzuki T., 1995, PhR, 256, 173

Janka H.-T., 2012, ARNPS, 62, 407

Jha S., Garnavich P., Challis P., Kirshner R., Berlind P., 1999, IAUC, 7339, 2

Jha S., Riess A.G., Kirshner R.P., 2007, ApJ, 659, 122

Kajava J., Fynbo J., Della Monica Ferreira D., Michalowski M., Zafar T., Sollerman J., 2008, CBET, 1479, 2

Kawabata K.S. et al., 2002, ApJ, 580, 39

Krisciunas K. et al., 2003, AJ, 125, 166

Kinugasa K., Kawakita H., Yamaoka H., 2004, IAU Circ., 8456, 2

Kniazev A.Yu., Arkhyz N., Engels D., Pustilnik S.A., Ugryumov A.V., Masegosa J., 1998, IAUC, 6900, 1

Kochanek C. S., Khan R., Dai X., 2012, ApJ, 759, 20

Kulkarni S.R., Ofek E.O., Rau A., et al. 2007, Nat., 447, 458

Langer N., 2012, ARA&A, in press

Larsson J., Levan A.J., Davies M.B., Fruchter A.S., 2007, MNRAS, 376, 1285

Leloudas G. et al., 2011, A&A, 530A, 95L

Lennarz D., Altmann D., Wiebusch C., 2012, A&A, 538, 120

Leonard D.C. et al. 2002a, PASP, 114, 35

Leonard D.C. et al. 2002b, AJ, 124, 2490

Leonard D.C., Chornock R., Filippenko A.V., 2003, IAUC, 8144, 2

Leonard D.C., IAUC, 8579, 2

Levesque E. M., Stringfellow G. S., Ginsburg A. G., Bally J., Keeney B. A., 2012, arXiv, arXiv:1211.4577

Li W.D., Modjaz M., Treffers R.R., Filippenko A.V., Leonard D.C., 1998, IAUC, 6882, 1

Li W.D., et al., 2003, PASP, 115, 453

Li W., Yamaoka H., Itagaki K., 2004, CBET, 100, 1

Li W., Wang X., Dyk S.D.V., Cuillandre J.-C., Foley R. J., Filippenko A.V., 2007, ApJ, 661, 1013

Li W. et al., 2011, Nature, 480, 348

Li W., et al., 2011, MNRAS, 412, 1441

Madison D., Li W., 2007, CBET, 1034, 1

Maeda K., et al., 2007, ApJ, 658, L5

Marchili N et al., 2010, A&A, 507, 47

Massey P., 2003, ARA&A, 41, 15

Massey P., 2002, ApJSS, 141, 81

Mattila S., Greimel R., Gerardy C., Meikle W.P.S., 2005, IAU Circ. 8474, 1

Mattila S., Smartt S.J., Eldridge J.J., Maund J.R., Crockett R.M., Danziger I.J., 2008, ApJ, 688, L91

Matheson T., Berlind P., 2001, IAUC, 7748, 3

Matheson T., Jha S., Challis P., Kirshner R., Calkinas M., 2001, IAUC, 7563, 2

Matheson T., Jha S., Challis P., Kirshner R., 2001, IAUC, 7597, 3

Matheson T., Jha S., Challis P., Kirshner R., Berlind P., 2001, IAUC, 7756, 4

Matheson T., Jha S., Challis P., Kirshner R., Rines K., 2001, IAUC, 7759, 2

Matheson T., Jha S., Challis P., Kirshner R., Berlind P., 2002, IAUC, 7800, 4

Matheson T., Challis P., Kirshner R., Berlind P., 2003, IAUC, 8206, 2

Matheson T., Challis P., Kirshner R., Berlind P., 2004, IAUC, 8269, 2

Matheson T., Challis P., Kirshner R., Berlind P., Huchra J., 2004, IAUC, 8293, 2

Matheson T., Challis P., Kirshner R., Berlind P., 2004, IAUC, 8304, 4

Maund, J. R., Smartt, S. J., Kudritzki, R. P., Podsiadlowski, P., & Gilmore, G. F. 2004, Nat., 427, 129

Maund J.R., Smartt S.J., Schweizer F., 2005, ApJ, 630, L33

Maund J.R., Smartt S.J, 2005, MNRAS, 360, 288

Maund J.R. et al., 2006, MNRAS, 369, 390

Maund J.R., Wheeler J.C., Patat F., Wang L., Baade D., Höflich P.A., 2007, 671, 1944

Maund J.R. et al. 2011, ApJ, 739, 37L

Mauerhan J. C., et al., 2012, arXiv, arXiv:1209.6320

Mattila S., Fraser M., Smartt S. J., Meikle W. P. S., Romero-Canizales C., Crockett R. M., Stephens A., 2012, arXiv, arXiv:1207.188

Mattila S., et al., 2012, ApJ, 756, 111

Maxwell A.J., Graham M.L., Parker A., Sadavoy S., Pritchet C.J., Hsiao E.Y., Balam D.D., 2010, CBET, 2245, 2

Mazzali P. A., Iwamoto K., Nomoto K., 2000, ApJ, 545, 407

Mazzali et al. 2002, ApJ, 572, 61

Mazzali P. A., et al., 2006, Natur, 442, 1018

Mazzali et al. 2007, ApJ, 670, 592

Mazzali P.A., Maurer I., Valenti S., Kotak R., Hunter D., 2010, MNRAS, 408, 87

Meikle P., Lucy L., Smartt S.J., Leibundgut B., Lundqvist P., Ostensen R., 2002, IAU Circ., 7811, 2

Milisavljevic D, Thorstensen J., Fesen R., 2011, CBET, 2745, 3

Milisavljevic D., Fesen R., Soderberg A., Pickering T., Kotze P., 2011, CBET, 2902, 2

Misra K., Pooley D., Chandra P., Bhattacharya D., Ray A.K., Sagar R., Lewin, W.H.G., 2007, MNRAS, 381, 280

Misra K., Sahu D.K., Anupama G.C., Pandey K., 2008, MNRAS, 389, 706

Modjaz M., Kirshner R., Challis P., Falco E., 2005a, IAUC, 8461, 3

Modjaz M., Kirshner R., Challis P., Calkins M., Hutchins

R., 2005b, IAUC, 8553, 2
Monard L.A.G., 2004, IAU Circ., 2004, IAU Circ., 8454, 1
Monard L.A.G., 2011a, CBET, 2899, 1
Monard L.A.G. et al. 2011b, CBET, 2946, 1
Moore M., Li W., Filippenko A.V., Chornock R., Foley R.J., 2004, IAUC, 8286, 2
Moriya T., Tominaga N., Tanaka M., Nomoto K., Sauer D.N., Mazzali P.A., Maeda K., Suzuki T., 2010, ApJ, 719, 1445
Morrell N., 2008, CBET, 1398, 1
Morrell N., Prieto J., Botti I., 2010, CBET, 2355, 1
Morrell N., Botti I., Lopez P. 2010, CBET, 2480, 2
Morrell N., Stritzinger M., Ho L., 2011, CBET, 2667, 2
Murphy J.W., Jennings Z.G., Williams B. Dalcanton J.J., Dolphin A.E., 2011, ApJ, 742L, 4M
Nakano S, Hirose Y., Kushida R., Kushida Y., Li W, 2002, IAU Circ., 7810, 1
Nakano S., Kawabata K.S., Minamoto H., Yamanaka M., 2011, CEBT, 2785, 1
Nakamura T., Mazzali P. A., Nomoto K., Iwamoto K., 2001, ApJ, 550, 991
Narayan G., et al., 2011, ApJ, 731, L11
Navasardyan H., Benetti S., Harutyunyan A., Turatto M., 2007, CBET, 939, 1
Navasardyan H., Benetti S., Harutyunyan A., Agnoletto I., Bufano F., Cappellaro E., Turatto M., 2008, CBET, 1325, 1
Nevski V., Maksym A., Elenin L., Schwartz M., 2010, CBET, 2245
Nomoto K.I., Iwamoto K., Suzuki T., 1995, PhR, 256, 173
Nugent P.E. et al., 2011, CBET, 2792, 1
Oliveira A.S., Steiner J.E., Cieslinski D., 2003, MNRAS, 346, 963
Olivares F., Folatelli G., 2007, CBET, 1120, 1
Paragi Z. et al., 2010, Nature, 463, 516
Pastorello A., Giro E., Altavilla G., Benetti S., 2000, IAUC, 7481, 2
Pastorello A. et al. 2004, MNRAS, 347, 74
Pastorello A. et al., 2005, MNRAS, 360, 950
Pastorello A. et al. 2006, MNRAS, 370, 1752
Pastorello A., et al. 2007a, Nature, 447, 829
Pastorello A., et al., 2007b, MNRAS, 376, 1301
Pastorello A., et al. 2008a, MNRAS, 389, 113
Pastorello A., et al. 2008b, MNRAS, 389, 955
Pastorello A. et al. A&A, 500, 1013
Pastorello A., Botticella M.T., Trundle C., et al. 2010, MNRAS, 408, 181
Pastorello A., et al., 2012, arXiv, arXiv:1210.3568
Patat F., Maia M., 1998, IAUC, 6888, 1
Patat F., Baade D., Wang L., 2005, CBET, 450, 1
Pejcha O., Prieto J., 2009, CBET 2049, 2
Pereyra A., Magalhẽs A.M., Rodrigues C.V., Silva C.R., Campos R., Hickel G., Cieslinski D., 2006, A&A, 454, 827
Phillips M., Hamuy M., Roth M., Morrell N., 2003, IAUC, 8086, 2
Phillips M., Folatelli G., Contreras C., Morrell N., 2006, CBET, 729, 1
Podsiadlowski P., 1992, PASP, 104, 717
Podsiadlowski Ph, Joss P.C., Hsu J.J.L., 1992, ApJ, 391, 246
Podsiadlowski P., Mazzali P. A., Nomoto K., Lazzati D., Cappellaro E., 2004, ApJ, 607, L17
Poznanski D., Prochaska J. X., Bloom J. S., 2012, MNRAS, 426, 1465
Poznanski D., Ganeshalingam M., Silverman J.M., Filippenko A.V., 2011, MNRAS, 415, L81
Pozzo M. et al. 2006, MNRAS, 368, 1169
Prieto J., 2006, CBET, 731, 1
Prieto J.L., Morrell N., Duhalde O., Monard L.A.G., 2010, CBET, 2364, 1
Prieto J.L., Szczygiel D.M., Kochanek C.S., Stanek K.Z., Thompson T.A., Beacom J.F., Garnavich P.M., Woodward C.E., 2010, arXiv, 1007.0011
Prieto J. L., Brimacombe J., Drake A. J., Howerton S., 2012, arXiv, arXiv:1210.3347
Puckett T., Langoussis A., Garradd G.J., 2000, IAU Circ., 7530, 1
Quimby R., Gerardy C., Hoeflich P., Wheeler J.C., Shetrone M., Riley V., 2004, IAU Circ., 8446, 1
Raskin C., Scannapieco E., Rhoads J., Della Valle M., 2008, ApJ, 689, 358R
Roy R., et al., 2011, ApJ, 736, 76
Ruiz-Lapuente P. et al. 2002, IAUC, 8028, 3
Rupak R. et al. 2011, MNRAS, 414, 167
Ryder S.D., Murrowood C.E., Stathakis R.A., 2006, MNRAS, 369, 32
Saha A., Thim F., Tammann G.A., Reindl B., Sandage A., 2006, ApJS, 165, 108
Sahu D.K., Anupama G.C., Srividya S., Muneer S., 2006, MNRAS, 372, 1315
Sahu D.K., Arora S., Anto P., 2011, CBET, 2658, 2
Salvo M., Schmidt B., Keller S., 2004, IAUC, 8432, 2
Salvo M., Blackman J., Schmidt B., Bessell M., 2006, CBET, 557, 1
Sana H., et al., 2012, Sci, 337, 444
Sanader A., Hamann W.-R., Todt H., A&A, 540, 133
Saviane I., Hibbard J.E., Rich R.M., 2004, AJ, 127, 660
Saviane I. Momany Y., da Costa G.S., Rich R.M., Hibbard J.E., 2008, ApJ, 678, 179
Schlegel D.J., Finkbeiner D.P., Davis M., 1998, ApJ, 500, 525
Schmidt B.., Salvo M., 2005, IAUC, 8496, 2
Schweizer F. et al., 2008, ApJ, 136, 1482
Silverman J.M., Kandrashoff M.T., Filippenko A.V., 2009, CBET, 1969, 2
Silverman J.M., Cenko S.B., Filippenko A.V., 2011, CBET, 2671, 2
Silverman J.M., Cenko S.B., Miller A.A., Nugent P.E., & Filippenko A.V., 2012, Central Bureau Electronic Telegrams, 3052, 2
Simon J.D. et al., 2007, ApJ, 671, L25
Smartt S.J., Vreeswijk P.M., Ramirez-Ruiz E., Gilmore G.F., Meikle W.P.S., Ferguson A.M.N., Knapen J.H., 2002, ApJ, 572, 147
Smartt S.J., Maund J.R., Gilmore G.F., Tout C.A., Kilkenny D., Benetti S., 2003, MNRAS, 343, 735
Smartt S.J., Eldridge J.J., Crockett R.M., Maund J.R., 2009, MNRAS, 395, 1409
Smartt S. J., 2009, ARA&A, 47, 63
Smith N. et al. 2009, ApJ, 697, 49
Smith N., et al., 2010, AJ, 139, 1451
Smith N., Li W., Filippenko A.V., Chornock R., 2011, MNRAS, 412, 1522S
Smith N., Li W., Silverman J.M., Ganeshalingam M., Fil-

ippenko A.V., 2011, MNRAS, 415, 773
Soderberg A.M., Chevalier R.A., Kulkarni S.R., Frail D.A., 2006, ApJ, 651, 1005
Soderberg A.M., Brunthaler A., Nakara E., Chevalier R.A., Bietenholz M.F., 2010, ApJ, 725, 922
Stanishev V., Adamo A., Micheva G., 2009, CBET, 1746, 1
Stanishev, V., & Pursimo, T. 2012, Central Bureau Electronic Telegrams, 2974, 3
Steiner J. E., Diaz M. P., 1998, PASP, 110, 276S
Steiner J.E., Oliveira A.S., 2005, A&A, 444, 895
Stritzinger M., et al., 2009, ApJ, 696, 713
Stritzinger M., Folatelli G., 2009, CBET 1902.
Stritzinger M., Forster F., Folatelli G., Pignata G., Hamuy M., 2009, CBET 1664.
Stritzinger M., Phillips M.M., Folatelli G., Foley R.J., 2010, CBET 2191, 1
Stockdale C.J., Williams C.L., Weiler K.W., Panagia N., Sramek R.A., Van Dyk S.D., Kelley M.T., 2007, ApJ, 671, 689
Strizinger M., 2010, CBET, 2158, 1
Stritzinger M., Foley R.J., 2011, CBET, 2899, 2
Swift B., Li W.D., Filippenko A.V., 2001, IAU Circ. 7618, 1
Takada-Hidai M., Aoki W., Zhao G., 2002, PASJ, 54, 899
Tanaka M., Kawabata K.S., Maeda K., Hattori T., Nomoto K., 2008, ApJ, 689, 1191
Tarchi A., Neininger N., Greve A., Klein U., Garrington S.T., Muxlow T.W.B., Pedlar A., Glendenning B.E., 2000, A&A, 358, 95
Taubenberger S., Pastorello A., Benetti S., Aceituno J., 2005a, IAUC, 8474, 3
Taubenberger S., Pastorello A., Mazzali P.A., Witham A., Guijarro A., 2005b, CBET 305, 1
Tonry J.L., Blakeslee J.P., Ajhar E.A., Dressler A., 2000, ApJ, 530, 625
Tsvetkov D.Y., 2006, Peremennye Zvezdy, 26, 3
Tully R.B., Rizzi L., Shaya E.J., Courtois H.M., Makarov D.I., Jacobs B.A., 2009, AJ, 138, 323
Turatto M., Benetti S., Cappellaro E., 2003, in Hillebrandt W., Leibundgut B., eds, From Twilight to Highlight: The Physics of Supernovae. Springer-Verlag, Berlin, p. 200
Turatto M., Benetti S., Pastorello A., 2007, astro-ph/0706.1086
Umbriaco G., Pietrogrande T., di Mille F., Agnoletto I., Harutyunyan A., Benetti S., 2007, CBET, 1174, 1
Utrobin V.P., Chugai N.N. Pastorello A., 2007, A&A, 475, 973
Valentini G., et al., 2003, ApJ, 595, 779
Valenti S. et al. 2008, ApJ, 673, 155
Valenti S., et al., 2008, MNRAS, 383, 1485
Valenti S. et al. 2009, Nature, 459, 674
Valenti S. et al. 2011a, MNRAS, 416, 3138
Valenti S., Benetti S., 2011b, CBET, 2749, 2
Valenti S., et al., 2011, MNRAS, 416, 3138
Valenti S., et al., 2012, ApJ, 749, L28
Vanbeveren D., Van Bever J., Belkus H., 2007, ApJ...662L.107V
Van Dyk S.D., Li W., Filippenko A.V., 2003c, PASP, 115, 1
Van Dyk S.D., Li W., Filippenko A.V., 2003a, PASP, 115, 1289
Van Dyk S.D., Li W., Filippenko A.V., 2003b, PASP, 115, 448
Van Dyk S.D. et al., 2011, ApJ, 741, 28L
Van Dyk, S. D., Cenko, S. B., Poznanski, D., et al. 2012, ApJ, 756, 131
Van Dyk S.D., Cenko S.B., SilvermanJ.M., Silverman J.M., Miller A.A., Filippenko A.V., Bloom J.S., 2012, The Astronomer's Telegram, 3971, 1
Vinkò J. et al 2006, MNRAS, 369, 1780
Vinkò J.¡ Marion G.H., Wheeler J.C., Foley R.J., Kirshner R.P., Rostopchin S., 2010, CBET, 2300, 1
Voss R., Nelemans G., 2008, Nature, 451, 802
Wei J.Y., Cao L., Qiu Y.L., Qiao Q.Y., Hu J.Y., Gu Q.S., 1999, IAUC,1 7124, 1
Wagner R. et al. 2004, PASP, 116, 326
Walborn N.R., Lasker B.M., Laidler V.G., Chu, Y.-H., 1987, ApJ, 321L, 41W
Wang L., Baade D., Höflich P., Wheeler J.C., 2003, ApJ, 592, 457
Wang L., Baade D., Clocchiatti A., Maund J., Patat F., Quinn J., 2008a, CBET, 1509, 1
Wang X. et al. 2008b, ApJ, 675, 626
Whitmore B.C., Zhang Q., Leitherer C., Fall S.M., Schweizer F., Miller B.W., 1999, AJ, 118, 4
Wood-Vasey W.M., Aldering G., Nugent P., Beutler B., Martin P., Papenkova M., Filippenko A.V., Hamuy M., 2002, IAUC, 799. 1
Xu D.W., Qiu Y.L., 2011, IAU Circ., 7555, 2
Yamanaka M., Arai A., Sakimoto K., Okushima T., Kawabata K.S., 2010, CBET, 2384, 1
Yoon S.-C., Langer N., Norman C., 2006, A&A, 460, 199Y
Yoon S.-C., Cantiello M., 2010, ApJ, 717L, 62
Yoon S.-C., Woosley S.E., Langer N., 2010, ApJ, 725, 940Y
Yoon S.-C., Grfener G., Vink J.S., Kozyreva A., Izzard R.G., 2012, A&A, 544, 11
Zhang T., Wang X., Li W., Zhou X., Ma J., Jiang Z., Chen J., 2006, AJ, 131, 2245
Zhang T., Zhou Z., 2011, 2708, 3

This paper has been typeset from a TEX/ LATEX file prepared by the author.

APPENDIX A: COMPLETE LIST OF SNE EMPLOYED IN THE SURVEY FOR PROGENITORS

Table A1: Core collapse SNe discovered between 1998-2012.25 in galaxies with recessional velocities less than 2000 kms^{-1} . The HST FOV column notes if the host galaxy has been observed by HST prior to explosion and if the position of the SN is "in" or "out" of the camera field-of-view.

Supernova	Galaxy	V_{vir}	Type	HST FOV	Comments
Core-collapse					
1998A	IC2627	1974	IIpec	...	Pastorello et al. (2005)SN1987A-like
1998S	NGC3877	1115	IIn	...	Fassia et al. (2001)
1998bm	IC2458	1810	II	...	Li et al. (1998), unknown subtype
1998bv	PGC2302994	1794	II-P	...	Kniazeb (1998)
1998dl	NGC1084	1298	II-P	...	Filippenko & De Breuck (1998a)
1998dn	NGC 337A	1004	II	...	Gómez & López (2000), unknown subtype
1999B	UGC7189	1962	II	...	Filippenko, Leonard & Riess (1998),unknown subtype
1999an	IC755	1605	II	in	Wei et al. (1999), unknown subtype
1999bg	IC758	1537	II-P	...	Ayani & Yamaoka (1999), amateur LC
1999br	NGC4900	1013	II-P	in	Pastorello et al. (2004) Low luminosity II-P
1999el	NGC6951	1704	IIn	out	Di Carlo et al. (2002)
1999em	NGC1637	615	II-P	out	Leonard et al. (2002a)
1999eu	NGC1097	1066	II-P	out	Pastorello et al. (2004) Low luminosity II-P
1999ev	NGC4274	1089	II-P	in	Garnavich et al. (1999), amateur LC.
1999ga	NGC2442	1168	II-L	...	Peculiar II-L Pastorello et al. (2009)
1999gi	NGC3184	765	II-P	in	Leonard et al. (2002b)
1999gn	NGC4303	1616	II-P	...	Pastorello et al. (2004) possible low luminosity II-P
1999gq	NGC4523	364	II-P	...	Jha et al. (1999), amateur LC
2000db	NGC3949	1020	II-P	out	Pastorello et al. (2000), amateur LC
2000ds	NGC2768	1622	Ib	in	Filippenko & Chornock (2000)
2000ew	NGC3810	1061	Ic	in	Gerardy et al. (2002)
2001B	IC391	1816	Ib	in	Chornock & Filippenko (2001), Tsvetkov Tsvetkov (2006)
2001X	NGC5921	1582	II-P	...	Tsvetkov (2006)
2001ci	NGC3079	1343	Ic	in	Filippenko & Chornock (2001)
2001du	NGC1365	1408	II-P	in	Smartt et al. (2003), Van Dyk, Li & Filippenko (2003b)
2001fv	NGC3512	1504	II-P	...	Matheson et al. (2001), amateur LC
2001fz	NGC2280	1720	II-P	...	Matheson er al. (2001), amateur LC
2001gd	NGC5033	1081	IIb	out	Stockdale et al. (2007)
2001ig	NGC7424	754	IIb	out	Ryder, Murrowood & Stathakis (2006), Maund et al. (2007)
2002E	NGC4129	1152	II	...	Matheson er al. (2002), unknown subtype
2002ao	UGC9299	1607	Ic	...	Pastorello et al. (2008a): narrow He lines[18]
2002ap	NGC628	686	Ic	out	Mazzali et al. (2002)
2002bu	NGC4242	737	IIn	out	Ayani, Kawabata & Yamaoka (2002), LC appears like II-L
2002hc	NGC2559	1370	II-L	...	Wood-Vasey et al. (2002), similar to 1979C
2002hh	NGC6946	318	II-P	out	Pozzo et al. (2006)
2002ji	NGC3655	1538	Ibc	...	Ruiz-Lapuente et al. (2002)
2002jz	UGC 2984	1527	Ic	...	Hamuy, Roth & Morrell (2002), somewhat uncertain
2003B	NGC1097	1066	II-P	...	M. Hamuy (priv. communication)
2003J	NGC4157	1011	II-P	...	Ayani, Hashimoto & Yamaoka (2003), amateur LC
2003Z	NGC2742	1518	II-P	...	Utrobin, Chugai & Pastorello (2007), low luminosity II-P
2003bg	MCG-05-10-15	1151	Ic	...	Soderberg et al. (2006), evidence of H during evolution
2003bk	NGC4316	1325	II	...	Phillips et al. (2003), subtype unknown
2003ed	NGC5303	1631	IIb	...	Leonard, Chornock & Filippenko (2003)
2003gd	NGC628	686	II-P	in	Hendry et al. (2005)
2003hn	NGC1448	911	II-P	out	M. Hamuy (priv. communication)
2003ie	NGC4051	917	II-P	...	Harutyunyan et al. (2008)

continued on next page

---

[18] SNe 2002ao and 2006jc have been termed Ibn as they show narrow He lines due to circumstellar He rich shells. We adopt Ic rather than Ibn as that more accurately reflects the nature of the event.

Table A1 continued

Supernova	Galaxy	V_{vir}	Type	HST flag	Comments
2003jg	NGC2997	914	Ibc	in	Howell (2003)
2004A	NGC6207	1090	II-P	in	Hendry et al. (2006)
2004C	NGC3683	1944	Ic	...	Matheson er al. (2004)
2004G	NGC5668	1673	II	out	Elias-Rosa et al. (2004a), subtype unknown
2004am	NGC3034	487	II-P	in	Mattila in prep.
2004ao	UGC10862	1812	Ib	...	Matheson et al. (2004)
2004bm	NGC3437	1381	Ic	...	Foley et al. (2004)
2004cm	NGC5486	1648	II-P	...	Connoly (2004), probable low-luminosity II-P
2004cz	ESO407-G09	1398	II-P	...	Foley, Pugh & Filippenko (2004)
2004dg	NGC5806	1439	II-P	in	Elias-Rosa et al. (2004b)
2004dj	NGC2403	370	II-P	out	Vinkò et al. (2006), Zhang et al. (2006)
2004dk	NGC6118	1645	Ib	...	Filippenko et al. (2004)
2004ep	IC2152	1683	II	...	Filippenko et al. (2004), subtype unknown
2004et	NGC6946	318	II-P	...	Misra et al. (2007), Sahu et al. (2006)
2004ez	NGC3430	1731	II-P	...	Filippenko et al. (2004), amateur LC
2004fc	NGC701	1727	II-P	...	Salvo, Schmidt & Keller (2004)
2004gk	IC3311	-53	Ic	...	Quimby et al. (2004)
2004gn	NGC4527	1780	Ic	out	1990B-like, SNIFS spectrum [19]
2004gq	NGC1832	1782	Ibc	-	Modjaz et al. (2005a), Gal-Yam et al. (2007)
2004gt	NGC4038	1559	Ibc	in	Gal-Yam et al. (2005),Maund, Smartt & Schweizer (2005)
2005V	NGC2146	1157	Ibc	in	Taubenberger et al. (2005a)
2005ad	NGC941	1535	II-P	...	M. Hamuy (priv. communication), probable II-P
2005ae	E209-G09	862	IIb	in	Filippenko & Foley (2005)
2005af	NGC4945	376	II-P	...	. Pereyra et al. (2006)
2005at	NGC6744	632	Ic	out	Schmidt & Salvo (2005)
2005ay	NGC3938	1017	II-P	...	Gal-Yam et al. (2008)
2005cs	NGC5194	702	II-P	in	Pastorello et al. (2006), low-luminosity II-P
2005cz	NGC4589	2283	Ib	in	Leonard (2005) [20]
2005kl	NGC4369	1248	Ic	...	Taubenberger et al. (2005b)
2006bc	NGC2397	1064	II-P	in	Patat, Baade & Wang (2006), amateur LC
2006bp	NGC3953	1285	II-P	...	Dessart et al. (2008)
2006jc	UGC4904	1830	Ic	..	Pastorello et al. (2007a), Foley et al. (2007): narrow He[21]
2006my	NGC4651	912.1	II-P	in	Li et al. (2007)
2006ov	NGC4303	1616	II-P	in	Li et al. (2007)
2007C	NGC4981	1687	Ib	...	Blondin & Calkins (2007a)
2007Y	NGC1187	1214	Ibc	...	Folatelli et al. (2007)
2007aa	NGC4030	1476	II-P	in	Folatelli, Gonzalez & Morrell (2007), our own LC
2007av	NGC3279	1436	II-P	...	Harutyunyan et al. (2007a), amateur LC
2007gr	NGC1058	634	Ic	in	Valenti et al. (2008a)
2007it	NGC5530	1045	II	...	Contreras et al. (2007), unknown subtype
2007oc	NGC7418A	1938	II-P	...	Olivares & Folatelli (2007), 1999em like
2007od	UGC12846	1810	II-P	...	Blondin & Calkins (2007b), 1999em like
2008M	E121-G26	1981	II-P	...	Folatelli, Olivares & Morrell (2008)
2008ax	NGC4490	797	IIb	in	Pastorello et al. (2008b)
2008bk	NGC7793	60	II-P	out	Mattila et al. (2008)
2008bo	NGC6643	1778	Ib	...	Navasardyan et al. (2008)
2008fb	UGC2813	1622	II	...	Kajava et al. (2008)
2008gz	NGC3672	1822	II-P	...	Rupak et al. (2011)
2008ij	NGC6643	1778	II	out	Challis (2008)
2008in	NGC4303	1616	II-P	out	Rou et al. (2011) low luminosity II-P
2009G	IC4444	1812	IIP	in	Stritzinger et al. (2009)
2009H	NGC1084	1298	IIP faint	in	Benetti et al. (2009)
2009N	NGC4487	1026	IIP faint	in	Challis (2009)
2009at	NGC5301	1749	II	...	Harutyunyan, Bufano & Benetti (2009)
2009bw	UGC2890	1404	II	...	Stanishev (2009)

continued on next page

---

[19] http://www.supernovae.net/sn2004/sn2004gn.jpg

[20] Although V_{vir} outside limit, the TF/SBF distance from Tonry et al. (2000) puts it within our distance limit and so it is included.

[21] SNe 2002ao and 2006jc have been termed Ibn as they show narrow He lines due to circumstellar He rich shells. We adopt Ic rather than Ibn as that more accurately reflects the nature of the event.

Table A1 continued

Supernova	Galaxy	V_{vir}	Type	HST flag	Comments
2009dd	NGC4088	989	II	in	Elias-Rosa et al. (2009b)
2009dq	IC2554	1108	II	...	Anderson et al. (2009)
2009em	NGC157	1600	Ic	out	Folatelli & Morrell (2009)
2009gj	NGC134	1404	IIb	out	Foley (2009)
2009hd	NGC3627	788	II	in	Elias-Rosa et al. (2011)
2009ib	NGC1559	1005	IIP	in	Stritzinger & Folatelli (2009)
2009js	NGC918	1536	IIP faint	out	Silverman, Kandrashoff & Filippenko (2009)
2009kr	NGC1832	1780	IIP/L	in	Fraser et al. (2010b), Elias-Rosa et al. (2010)
2009ls	NGC3423	1033	IIL	in	Pejcha & Prieto (2009)
2009md	NGC3389	1356	II	in	Fraser et al. (2011)
2009mg	ESO121-G26	1981	IIb	out	Strizinger (2010)
2009mk	ESO293-34	1305	IIb	...	Chornock & Berger (2009)
2010br	NGC4051	917	Ibc	in	Maxwell et al. (2010)
2010gi	IC4660	1538	IIb	...	Yamanaka et al. (2010)
2011am	NGC4219	1809	Ib	in	Morrell, Stritzinger & Ho (2011)
2011aq	NGC1056	1620	II	...	Silverman, Cenko & Filippenko (2011)
2011dh	NGC5194	702	IIb	in	Maund et al. (2011), Van Dyk et al. (2011)
2011dq	NGC337	1572	IIP faint		Valenti & Benetti (2011b)
2011hp	NGC4219	1811	Ic	in	Monard (2011a)
2011hs	IC5267	1524	IIb	out	Milisavljevic et al. (2011)
2011ja	NGC4945	376	IIP	out	Monard et al. (2011b)
2011jm	NGC4809	979	Ic	...	Howerton et al. (2011)
2012A	NGC3239	820	II	...	Stanishev & Pursimo (2012)
2012P	NGC5806	1439	IIb	in	Borsato et al. (2012)
2012au	NGC4790	1344	Ib	in	Silverman et al. (2012)
2012aw	NGC3351	826	IIP	in	Fraser et al. (2012), Van Dyk et al. (2012)
Type Ia					
1998aq	NGC3982	1388	Ia	in	Branch (2003)
1998bn	NGC4462	1718	Ia	...	Patat & Maia (1998)
1998bu	NGC3368	941	Ia	out	Hernandez et al. (2000)
1998dm	PGC005341	1882	Ia	...	Filippenko & De Breuck (1998b)
1999by	NGC2841	831	Ia	...	Garnavich et al. (2004)
2000E	NGC6951	1704	Ia	...	Valentini et al. (2003)
2001dp	NGC3953	1285	Ia	out	Ayani et al. (2001)
2001el	NGC1448	911	Ia	out	Krisciunas et al. (2003)
2001fu	M-03-23-11	1564	Ia	...	Matheson & Berlind (2001)
2002bo	NGC3190	1410	Ia	out	Benetti et al. (2004)
2002cv	NGC3190	1410	Ia	out	Elias-Rosa et al. (2008)
2002fk	NGC1309	1986	Ia	...	Ayani & Yamoka (2002)
2003cg	NGC3169	1237	Ia	in	Elias-Rosa et al. (2006)
2003gs	NGC936	1279	Ia	in	Gonzalez, Morrell & Hamuy (2003)
2003hv	NGC1201	1490	Ia	in	Gerardy et al. (2007)
2003hx	NGC2076	1974	Ia	...	Misra et al. (2008)
2003if	NGC1302	1506	Ia	...	Matheson er al. (2003)
2004W	NGC4649	1198	Ia	in	Moore et al. (2004)
2004ab	NGC5054	1708	Ia	in	Matheson er al. (2004)
2004ea	M-03-11-19	1792	Ia	...	Gonzalez et al. (2004)
2005cf	M-01-39-03	1977	Ia	...	Pastorello et al. (2007b)
2005cn	NGC5061	1944	Ia	out	Modjaz et al. (2005b)
2005df	NGC1559	1005	Ia	in	Gerardy et al. (2007)
2005ke	NGC1371	1272	Ia	...	Immler et al. (2006)
2006E	NGC5338	900	Ia	...	Aldering et al. (2006)
2006X	NGC4321	1677	Ia	in	Wang et al. (2008)
2006ce	NGC908	1344	Ia	...	Blackmass, Schmidt & Kerzendorf (2006)
2006dd	NGC1316	1556	Ia	in	Salvo et al. (2006)
2006mq	ESO494-G26	779.4	Ia	..	Preito (2006)
2006mr	NGC1316	1556	Ia	in	Phillips et al. (2006)
2007af	NGC5584	1704	Ia	in	Simon et al. (2007)
2007bm	NGC3672	1824	Ia	...	Navasardyan et al. (2007)
2007gi	NGC4036	1629	Ia	in	Harutyunyan et al. (2007b)
2007le	NGC7721	1971	Ia	...	Filippenko et al. (2007)
2007on	NGC1404	1717	Ia	in	Voss & Nelemans (2008)
2007sa	NGC3499	1761	Ia	...	Agnoletto et al. (2007)
2007sr	NGC4038	1559	Ia	in	Umbriaco et al. (2007)

continued on next page

Table A1 continued

Supernova	Galaxy	V_{vir}	Type	HST flag	Comments
2008cq	PGC731537	1191	Ia	...	Morrell (2008)
2008fp	ESO428-G14	1504	Ia	in	Wang et al. (2008a)
2008ge	NGC1527	910	Ia	in	Foley et al. (2010b) 2002cx-like
2008ha	UGC12682	1469	Ia	...	Valenti et al. (2009), Foley et al. (2009) 2002cx-like
2010ae	ESO162-17	818	Ia	...	Strizinger et al. (2010), 2002cx-like
2010el	NGC1566	1221	Ia	in	Bessel & Schmidt (2010), 2002cx-like
2010ey	ESO548-76	1312	Ia	...	Morrell & Prieto (2010)
2010fz	NGC2967	1861	Ia	...	Preito et al. (2010)
2010ih	NGC2325	1954	Ia	out	Morrell, Botti & Lopez (2010)
2011B	NGC2655	1653	Ia	out	Bakanm Graham & Hsiao (2011)
2011ae	MCG-03-30-19	1747	Ia	...	Sahu, Arora & Anto (2011)
2011at	MCG-02-24-027	1922	Ia	...	Bufano, Benetti & Pastorello (2011)
2011by	NGC3972	1086	Ia	in	Zhang & Zhou (2011)
2011dm	UGC11861	1756	Ia	...	Milisavljevic, Thorstensen & Fesen (2011)
2011ek	NGC918	1536	Ia	out	Nakano et al. (2011)
2011fe	M101	502	Ia	in	Nugent et al. (2011), Li et al. (2011)
2011iv	NGC1404	1717	Ia	in	Foley et al. (2012)
2011iy	NGC4984	1180	Ia	...	Chen et al. (2011)
Unclassified					
1998cf	NGC3504	1669	...	in	
1999gs	NGC4725	1206	...	out	
2008eh	NC2997	91	...	out	
2008iz	NGC3034	487	CC	in	
Uncertain or non-supernova					
1999bw	NGC3198	850.9	LBV	out	Filippenko, Li & Modjaz (1999)[22]
2000ch	NGC3432	778.6	LBV	out	Wagner et al. (2004)
2001ac	NGC3504	1668.9	LBV	out	Matheson et al. (2001)
2002kg	NGC2403	370	LBV	out	Maund et al. (2006)
2003gm	NGC5334	1433.8	LBV	in	Maund et al. (2006)
2007sv	UGC5979	1377.0	LBV	...	Harutyunyan et al. (2007c), our own data
2006fp	UGC12182	1807.5	LBV	..	Blondin et al. (2006)
M85 OT2006-1	M85	957	SN or OT?	in	Kulkarni et al. (2007)
2008S	NGC6946	318	IIn or OT?	out	Botticella et al. (2009)
NGC300 OT2008-1	NGC300	100	SN or OT?	in	Pastorello et al. (2010)
2009ip	NGC7259	1596	LBV	in	Pastorello et al. (2012) and references within
2010U	NGC4214	479	Nova	in	Humphreys et al. (2010)
2010da	NGC300	-38	LBV/08S-like	in	Binder et al. (2011)
2010dn	NGC3184	755	LBV/08S-like	in	Vinkò et al. (2010)
2011ht	UGC 5460		IIn		IMPOSTOR

---

[22] http://etacar.umn.edu/etainfo/related/